arXiv:1112.3792v1 [math.RT] 16 Dec 2011

A New Functor from D_5-Mod to E_6-Mod [1]

Xiaoping Xu

Hua Loo-Keng Key Mathematical Laboratory

Institute of Mathematics, Academy of Mathematics & System Sciences

Chinese Academy of Sciences, Beijing 100190, P.R. China [2]

Abstract

We find a new representation of the simple Lie algebra of type E_6 on the polynomial algebra in 16 variables, which gives a fractional representation of the corresponding Lie group on 16-dimensional space. Using this representation and Shen's idea of mixed product, we construct a functor from D_5-**Mod** to E_6-**Mod**. A condition for the functor to map a finite-dimensional irreducible D_5-module to an infinite-dimensional irreducible E_6-module is obtained. Our general frame also gives a direct polynomial extension from irreducible D_5-modules to irreducible E_6-modules. The obtained infinite-dimensional irreducible E_6-modules are $(\mathcal{G}, K)$-modules in terms of Lie group representations. The results could be used in studying the quantum field theory with E_6 symmetry and symmetry of partial differential equations.

1 Introduction

A quantum field is an operator-valued function on a certain Hilbert space, which is often a direct sum of infinite-dimensional irreducible modules of a certain Lie algebra (group). The Lie algebra of two-dimensional conformal group is exactly the Virasoro algebra. The minimal models of two-dimensional conformal field theory were constructed from direct sums of certain infinite-dimensional irreducible modules of the Virasoro algebra, where a distinguished module called, the *vacuum module*, gives rise to a vertex operator algebra.

It is well known that n-dimensional projective group gives rise to a non-homogenous representation of the Lie algebra $sl(n+1,\mathbb{C})$ on the polynomial functions of the projective space. Using Shen's mixed product for Witt algebras, Zhao and the author [ZX] generalized the above representation of $sl(n+1,\mathbb{C})$ to a non-homogenous representation on the tensor space of any finite-dimensional irreducible $gl(n,\mathbb{C})$-module with the polynomial space. Moreover, the structure of such a representation was completely determined by employing projection operator techniques (cf. [Gm]) and the well-known Kostant's characteristic identities (cf. [K]). The result can be used to study the quantum field theory with $sl(n+1,\mathbb{C})$ as the symmetry. Furthermore, we [XZ] generalize the conformal representation of of $o(n+2,\mathbb{C})$ to a non-homogenous representation of $o(n+2,\mathbb{C})$ on the tensor space of any finite-dimensional irreducible $o(n,\mathbb{C})$-module with a polynomial space by Shen's idea of mixed product for Witt algebras. It turns out that a hidden central transformation is involved. More importantly, we find a condition on the constant value taken by the central transformation such that the generalized conformal representation is irreducible. The result would be useful in higher-dimensional conformal field theory.

---

[1] 2000 Mathematical Subject Classification. Primary 17B10, 17B25;Secondary 22E46.

[2] Research supported by China NSF 11171324

This paper is the third work in the program of studying quantum-field motivated representations of finite-dimensional simple Lie algebras. It is well known that the minimal dimension of irreducible modules over the simple Lie algebra of type E_6 is 27. Based on a grading of the simple Lie algebra of type E_6, we find a first-order differential operator representation of the Lie algebra on the polynomial algebra in 16 independent variables. In fact, the corresponding Lie group representation is given by fractional transformations on 16-dimensional space. Using this representation and Shen's idea of mixed product, we construct a new functor from D_5-**Mod** to E_6-**Mod**, where a hidden central transformation is involved. More importantly, a condition for the functor to map a finite-dimensional irreducible D_5-module to an infinite-dimensional irreducible E_6-module in terms of the constant value taken by the central transformation is obtained. Our general frame also gives a direct polynomial extension from irreducible D_5-modules to irreducible E_6-modules, which can be applied to obtain explicit bases of irreducible E_6-modules from those of irreducible D_5-modules. The result could be useful in understanding the quantum field theory with E_6 symmetry. Our fractional representation of the E_6 Lie group could also be used in symmetry analysis of partial differential equations just as the conformal representation of orthogonal Lie groups do. Our infinite-dimensional irreducible E_6-modules are $(\mathcal{G},K)$-modules in terms of the corresponding Lie group representations.

The E_6 Lie algebra and group are popular mathematical objects with broad applications. Dickson [D] (1901) first realized that there exists an E_6-invariant trilinear form on its 27-dimensional basic irreducible module. The 78-dimensional simple Lie algebra of type E_6 can be realized by all the derivations and multiplication operators with trace zero on the 27-dimensional exceptional simple Jordan algebra (e.g., cf. [T], [Ad]). Aschbacher [As] used the Dickson form to study the subgroup structure of the group E_6. Bion-Nadal [B-N] proved that the E_6 Coxeter graph can be realized as a principal graph of subfactor of the hyperfinite Π_1 factor. Brylinski and Kostant [BK] obtained a generalized Capelli identity on the minimal representation of E_6. Binegar and Zierau [BZ] found a singular representation of E_6. Ginzburg [G] proved that the twisted partial L-function on the

27-dimensional representation of $GE_6(\mathbb{C})$ is entire except the points 0 and 1. Iltyakov [I] showed that the field of invariant rational functions of E_6 on the direct sum of finite copies of the basic module and its dual is purely transcendental. Suzuki and Wakui [SW] studied the Turaev-Viro-Ocneanu invariant of 3-manifilds derived from the E_6-subfactor. Moreover, Cerchiai and Scotti [CS] investigated the mapping geometry of the E_6 group. Furthermore, the (A_2, G_2) duality in E_6 was obtained by Rubenthaler [R]. In [X2], the author proved that the space of homogeneous polynomial solutions with degree m for the dual cubic Dickson invariant differential operator is exactly a direct sum of $[\![m/2]\!]+1$ explicitly determined irreducible E_6-submodules and the whole polynomial algebra is a free module over the polynomial algebra in the Dickson invariant generated by these solutions. Moreover, we found in [X3] that the weight matrices of E_6 on its minimal irreducible modules and adjoint modules all generate ternary orthogonal codes with large minimal distances.

Okamoto and Marshak [OM] constructed a grand unification preson model with E_6 metacolor. The E_6 Lie algebra was used in [HH] to explain the degeneracies encountered in the genetic code as the result of a sequence of symmetry breakings that have occurred during its evolution. Wang [W] identified Geoner's model with twisted LG model and E_6 singlets. Morrison, Pieruschka and Wybourne [MPW] constructed the E_6 interacting boson model. Berglund, Candelas et al. [BCDH] studied instanton contributions to the masses and couplings of E_6 singles. Haba and Matsuoka [HM] found large lepton flavor mixing in the E_6-type unification models. Ghezelbash, Shafiekhani and Abolbasani [GSA] derived explicitly a set of Picard-Fuchs equations of $N=2$ supersymmetric E_6 Yang-Mills theory. Anderson and Bla*ž*ek [AB1-AB3] found certain Clebsch-Gordan coefficients in connection with the E_6 unification model building. Fernández-Nú*ã*ez, Garcia-Fuertes and Perelomov [FGP] used the quantum Calogero-Sutherland model corresponding to the root system of E_6 to calculate Clebach-Gordan series for this algebra. Howl and King [HK] proposed a minimal E_6 supersymmetric standard model which allows Planck scale unification, provides a solution to the μ problem and predicts a new Z'. Das and Laperashvili [DL] studied Preon model related to family replicated E_6 unification.

This work further reveals new beauties of the simple Lie algebra of type E_6. In Section 2, we construct the spin representation of $o(10,\mathbb{C})$ in terms of first-order differential operators on the polynomial algebra in 16 independent variables from the lattice-construction of the simple Lie algebra of type E_6. We determine the decomposition of the polynomial algebra into irreducible $o(10,\mathbb{C})$-submodules in Section 3 by means of partial differential equations. In Section 4, we realized the simple Lie algebra of type E_6 in terms of first-order differential operators on the polynomial algebra in 16 independent variables.

Section 5 is devoted to the explicit presentation of the functor from D_5-**Mod** to E_6-**Mod**. Finally in Section 6, we determine a condition for the functor to map a finite-dimensional irreducible D_5-module to an infinite-dimensional irreducible E_6-module.

2 Polynomial Representation of $o(10,\mathbb{C})$ via E_6

We start with the root lattice construction of the simple Lie algebra of type E_6. As we all know, the Dynkin diagram of E_6 is as follows:



For convenience, we will use the notion $\overline{i,i+j}=\{i,i+1,i+2,...,i+j\}$ for integer i and positive integer j throughout this paper. Let $\{\alpha_i\mid i\in\overline{1,6}\}$ be the simple positive roots corresponding to the vertices in the diagram, and let Φ_{E_6} be the root system of E_6. Set

$$Q_{E_6}=\sum_{i=1}^{6}\mathbb{Z}\alpha_i, \tag{2.1}$$

the root lattice of type E_6. Denote by $(\cdot,\cdot)$ the symmetric $\mathbb{Z}$-bilinear form on Q_{E_6} such that

$$\Phi_{E_6}=\{\alpha\in Q_{E_6}\mid(\alpha,\alpha)=2\}. \tag{2.2}$$

Define a map $F:Q_{E_6}\times Q_{E_6}\to\{1,-1\}$ by

$$F(\sum_{i=1}^{6}k_i\alpha_i,\sum_{j=1}^{6}l_j\alpha_j)=(-1)^{\sum_{i=1}^{6}k_il_i+k_1l_3+k_4l_2+k_3l_4+k_5l_4+k_6l_5},\qquad k_i,l_j\in\mathbb{Z}. \tag{2.3}$$

Then for $\alpha,\beta,\gamma\in Q_{E_6}$,

$$F(\alpha+\beta,\gamma)=F(\alpha,\gamma)F(\beta,\gamma),\;\;F(\alpha,\beta+\gamma)=F(\alpha,\beta)F(\alpha,\gamma), \tag{2.4}$$

$$F(\alpha,\beta)F(\beta,\alpha)^{-1}=(-1)^{(\alpha,\beta)},\;\;F(\alpha,\alpha)=(-1)^{(\alpha,\alpha)/2}. \tag{2.5}$$

In particular,

$$F(\alpha,\beta)=-F(\beta,\alpha)\qquad\text{if }\;\alpha,\beta,\alpha+\beta\in\Phi_{E_6}. \tag{2.6}$$

Denote

$$H=\bigoplus_{i=1}^{6}\mathbb{C}\alpha_i \tag{2.7}$$

and $\mathbb{C}$-bilinearly extend $(\cdot,\cdot)$ on H. Then the simple Lie algebra of type E_6 is

$$\mathcal{G}^{E_6}=H\oplus\bigoplus_{\alpha\in\Phi_{E_6}}\mathbb{C}E_\alpha \tag{2.8}$$

with the Lie bracket $[\cdot,\cdot]$ determined by:

$$[H,H]=0,\ \ [h,E_\alpha]=-[E_\alpha,h]=(h,\alpha)E_\alpha,\ \ [E_\alpha,E_{-\alpha}]=-\alpha, \tag{2.9}$$

$$[E_\alpha,E_\beta]=\begin{cases}0 & \text{if } \alpha+\beta\not\in\Phi_{E_6},\\ F(\alpha,\beta)E_{\alpha+\beta} & \text{if } \alpha+\beta\in\Phi_{E_6}\end{cases} \tag{2.10}$$

(e.g., cf. [Ka], [X1]). Moreover, we define a bilinear form $(\cdot|\cdot)$ on $\mathcal{G}^{E_6}$ by

$$(h_1|h_2)=(h_1,h_2),\ \ (h|E_\alpha)=0,\ \ (E_\alpha|E_\beta)=-\delta_{\alpha+\beta,0} \tag{2.11}$$

for $h_1,h_2\in H$ and $\alpha,\beta\in\Phi_{E_6}$. It can be verified that $(\cdot|\cdot)$ is a $\mathcal{G}^{E_6}$-invariant form, that is,

$$([u,v]|w)=(u|[v,w])\qquad\text{for }\ u,v\in\mathcal{G}^{E_6}. \tag{2.12}$$

Let

$$Q^{D_5}=\sum_{i=1}^5\mathbb{Z}\alpha_i,\qquad \Phi_{D_5}=\Phi_{E_6}\bigcap Q^{D_5}. \tag{2.13}$$

Then

$$\mathcal{G}^{D_5}=\sum_{i=1}^5\mathbb{C}\alpha_i+\sum_{\beta\in\Phi_{D_5}}\mathbb{C}E_\beta \tag{2.14}$$

forms a Lie subalgebra of $\mathcal{G}^{E_6}$, which is isomorphic to the orthogonal Lie algebra

$$\begin{aligned} o(10,\mathbb{C}) \ \ &= \sum_{1\leq p<q\leq 5}[\mathbb{C}(E_{p,n+q}-E_{q,n+p})+\mathbb{C}(E_{n+p,q}-E_{n+q,p})] \\ &\quad +\sum_{i,j=1}^5\mathbb{C}(E_{i,j}-E_{n+j,n+i}). \end{aligned} \tag{2.15}$$

Denote by $\Phi_{E_6}^+$ the set of positive roots of E_6 and by $\Phi_{D_5}^+$ the set of positive roots of D_5. We find the elements of $\Phi_{D_5}^+$:

$$\alpha_r\ (r\in\overline{1,5}),\ \alpha_1+\alpha_3,\ \alpha_2+\alpha_4,\ \alpha_3+\alpha_4,\ \alpha_4+\alpha_5,\ \alpha_1+\alpha_3+\alpha_4,\ \alpha_2+\alpha_3+\alpha_4, \tag{2.16}$$

$$\alpha_2+\alpha_4+\alpha_5,\ \alpha_3+\alpha_4+\alpha_5,\sum_{r=1}^4\alpha_r,\ \alpha_1+\alpha_3+\alpha_4+\alpha_5, \tag{2.17}$$

$$\sum_{i=2}^5\alpha_i,\ \sum_{s=1}^5\alpha_s,\ \alpha_4+\sum_{i=2}^5\alpha_i,\ \alpha_4+\sum_{i=1}^5\alpha_i,\ \alpha_3+\alpha_4+\sum_{i=1}^5\alpha_i. \tag{2.18}$$

Moreover, the elements in $\Phi_{E_6}^+\setminus\Phi_{D_5}^+$ are:

$$\alpha_6,\ \alpha_5+\alpha_6,\ \sum_{r=4}^6\alpha_r,\ \sum_{i=3}^6\alpha_i,\ \alpha_2+\sum_{r=4}^6\alpha_r,\ \sum_{i=2}^6\alpha_i,\ \alpha_1+\sum_{i=3}^6\alpha_i,\ \sum_{i=1}^6\alpha_i,\ \alpha_4+\sum_{i=2}^6\alpha_i, \tag{2.19}$$

$$\alpha_4+\sum_{i=1}^6\alpha_i,\ \alpha_4+\alpha_5+\sum_{i=2}^6\alpha_i,\ \alpha_3+\alpha_4+\sum_{i=1}^6\alpha_i,\ \alpha_4+\alpha_5+\sum_{i=1}^6\alpha_i, \tag{2.20}$$

$$\sum_{i=1}^{6}\alpha_i+\sum_{r=3}^{5}\alpha_r,\ \alpha_4+\sum_{i=1}^{6}\alpha_i+\sum_{r=3}^{5}\alpha_r,\ \alpha_4+\sum_{i=1}^{6}\alpha_i+\sum_{r=2}^{5}\alpha_r. \tag{2.21}$$

For convenience, we denote

$$\xi_1=E_{\alpha_6},\ \ \xi_2=E_{\alpha_5+\alpha_6},\ \ \xi_3=E_{\sum_{r=4}^6\alpha_r},\ \ \xi_4=E_{\sum_{i=3}^6\alpha_i}, \tag{2.22}$$

$$\xi_5=E_{\alpha_2+\sum_{r=4}^6\alpha_r},\ \ \xi_6=E_{\sum_{i=2}^6\alpha_i},\ \ \xi_7=E_{\alpha_1+\sum_{i=3}^6\alpha_i},\ \ \xi_8=E_{\sum_{i=1}^6\alpha_i}, \tag{2.23}$$

$$\xi_9=E_{\alpha_4+\sum_{i=2}^6\alpha_i},\ \ \xi_{10}=E_{\alpha_4+\sum_{i=1}^6\alpha_i},\ \xi_{11}=E_{\alpha_4+\alpha_5+\sum_{i=2}^6\alpha_i}, \tag{2.24}$$

$$\xi_{12}=E_{\alpha_3+\alpha_4+\sum_{i=1}^6\alpha_i},\ \ \xi_{13}=E_{\alpha_4+\alpha_5+\sum_{i=1}^6\alpha_i},\ \ \xi_{14}=E_{\sum_{i=1}^6\alpha_i+\sum_{r=3}^5\alpha_r}, \tag{2.25}$$

$$\xi_{15}=E_{\alpha_4+\sum_{i=1}^6\alpha_i+\sum_{r=3}^5\alpha_r},\ \ \xi_{16}=E_{\alpha_4+\sum_{i=1}^6\alpha_i+\sum_{r=2}^5\alpha_r}, \tag{2.26}$$

$$\eta_1=E_{-\alpha_6},\ \ \eta_2=E_{-\alpha_5-\alpha_6},\ \ \eta_3=E_{-\sum_{r=4}^6\alpha_r},\ \ \eta_4=E_{-\sum_{i=3}^6\alpha_i}, \tag{2.27}$$

$$\eta_5=E_{-\alpha_2-\sum_{r=4}^6\alpha_r},\ \ \eta_6=E_{-\sum_{i=2}^6\alpha_i},\ \ \eta_7=E_{-\alpha_1-\sum_{i=3}^6\alpha_i},\ \ \eta_8=E_{-\sum_{i=1}^6\alpha_i}, \tag{2.28}$$

$$\eta_9=E_{-\alpha_4-\sum_{i=2}^6\alpha_i},\ \ \eta_{10}=E_{-\alpha_4-\sum_{i=1}^6\alpha_i},\ \eta_{11}=E_{-\alpha_4-\alpha_5-\sum_{i=2}^6\alpha_i}, \tag{2.29}$$

$$\eta_{12}=E_{-\alpha_3-\alpha_4-\sum_{i=1}^6\alpha_i},\ \ \eta_{13}=E_{-\alpha_4-\alpha_5-\sum_{i=1}^6\alpha_i},\ \ \eta_{14}=E_{-\sum_{i=1}^6\alpha_i-\sum_{r=3}^5\alpha_r}, \tag{2.30}$$

$$\eta_{15}=E_{-\alpha_4-\sum_{i=1}^6\alpha_i-\sum_{r=3}^5\alpha_r},\ \ \eta_{16}=E_{-\alpha_4-\sum_{i=1}^6\alpha_i-\sum_{r=2}^5\alpha_r}. \tag{2.31}$$

Set

$$\mathcal{G}_+=\sum_{i=1}^{16}\mathbb{C}\xi_i,\qquad \mathcal{G}_-=\sum_{i=1}^{16}\mathbb{C}\eta_i,\qquad \mathcal{G}_0=\mathcal{G}^{D_5}+\mathbb{F}\alpha_6. \tag{2.32}$$

It is straightforward to verify that $\mathcal{G}_\pm$ are ableian Lie subalgebras of $\mathcal{G}^{E_6}$, $\mathcal{G}_0$ is a reductive Lie subalgebra of $\mathcal{G}^{E_6}$ and

$$\mathcal{G}^{E_6}=\mathcal{G}_-\oplus\mathcal{G}_0\oplus\mathcal{G}_+. \tag{2.33}$$

Moreover, $\mathcal{G}_\pm$ form irreducible $\mathcal{G}_0$-submodules with respect to the adjoint representation of $\mathcal{G}^{E_6}$. Furthermore,

$$(\xi_i|\eta_j)=-\delta_{i,j}\qquad \text{for}\ \ i,j\in\overline{1,16} \tag{2.34}$$

by (2.11). Expression (2.12) shows that $\mathcal{G}_+$ is isomorphic to the dual $\mathcal{G}_0$-module of $\mathcal{G}_-$.

Set

$$\mathcal{A}=\mathbb{C}[x_1,x_2,...,x_{16}], \tag{2.35}$$

the polynomial algebra in $x_1,x_2,...,x_{16}$. Write

$$[u,\eta_i]=\sum_{j=1}^{16}\varphi_{i,j}(u)\eta_j\qquad \text{for}\ \ i\in\overline{1,16},\ u\in\mathcal{G}_0, \tag{2.36}$$

where $\varphi_{i,j}(u)\in\mathbb{C}$. Define an action of $\mathcal{G}_0$ on $\mathcal{A}$ by

$$u(f)=\sum_{i,j=1}^{16}\varphi_{i,j}(u)x_j\partial_{x_i}(f)\qquad \text{for}\ \ u\in\mathcal{G}_0,\ f\in\mathcal{A}. \tag{2.37}$$

Then $\mathcal{A}$ forms a $\mathcal{G}_0$-module and the subspace

$$V=\sum_{i=1}^{16}\mathbb{C}x_i \tag{2.38}$$

forms a $\mathcal{G}_0$-submodule isomorphic to $\mathcal{G}_-$, where the isomorphism is determined by $x_i\mapsto\eta_i$ for $i\in\overline{1,16}$.

Denote by $\mathbb{N}$ the set of nonnegative integers. Write

$$x^\alpha=\prod_{i=1}^{16}x_i^{\alpha_i},\;\;\partial^\alpha=\prod_{i=1}^{16}\partial_{x_i}^{\alpha_i}\qquad\text{for }\;\alpha=(\alpha_1,\alpha_2,...,\alpha_{16})\in\mathbb{N}^{16}. \tag{2.39}$$

Let

$$\mathbb{A}=\sum_{\alpha\in\mathbb{N}^{16}}\mathcal{A}\partial^\alpha \tag{2.40}$$

be the algebra of differential operators on $\mathcal{A}$. Then the linear transformation τ determined by

$$\tau(x^\beta\partial^\gamma)=x^\gamma\partial^\beta\qquad\text{for }\;\beta,\gamma\in\mathbb{N}^{16} \tag{2.41}$$

is an involutive anti-automorphism of $\mathbb{A}$.

According to (2.9) and (2.10), we have the Lie algebra isomorphism $\nu:o(10,\mathbb{C})\rightarrow\mathcal{G}^{D_5}$ determined by the generators:

$$\nu(E_{1,2}-E_{7,6})=E_{\alpha_1},\;\;\nu(E_{2,3}-E_{8,7})=E_{\alpha_3},\;\;\nu(E_{3,4}-E_{9,8})=E_{\alpha_4}, \tag{2.42}$$

$$\nu(E_{4,5}-E_{10,9})=E_{\alpha_5},\;\;\nu(E_{4,10}-E_{5,9})=E_{\alpha_2},\;\;\nu(E_{2,1}-E_{6,7})=-E_{-\alpha_1} \tag{2.43}$$

$$\nu(E_{3,2}-E_{7,8})=-E_{-\alpha_3},\;\;\nu(E_{4,3}-E_{8,9})=-E_{-\alpha_4},\;\;\nu(E_{5,4}-E_{9,10})=-E_{-\alpha_5}, \tag{2.44}$$

$$\nu(E_{10,4}-E_{9,5})=-E_{-\alpha_2},\;\;\nu(E_{1,1}-E_{6,6})=\alpha_1+\alpha_3+\alpha_4+\frac{1}{2}(\alpha_2+\alpha_5), \tag{2.45}$$

$$\nu(E_{2,2}-E_{7,7})=\alpha_3+\alpha_4+\frac{1}{2}(\alpha_2+\alpha_5),\;\;\nu(E_{3,3}-E_{8,8})=\alpha_4+\frac{1}{2}(\alpha_2+\alpha_5), \tag{2.46}$$

$$\nu(E_{4,4}-E_{9,9})=\frac{1}{2}(\alpha_2+\alpha_5),\qquad\nu(E_{5,5}-E_{10,10})=\frac{1}{2}(\alpha_2-\alpha_5). \tag{2.47}$$

Then $\mathcal{A}$ becomes an $o(10,\mathbb{C})$-module with respect to the action

$$A(f)=\nu(A)(f)\qquad\text{for }\;A\in o(10,\mathbb{C}),\;f\in\mathcal{A}. \tag{2.48}$$

Thanks to (2.9), (2.10), (2.27)-(2.31), (2.36) and (2.37), we have

$$(E_{1,2}-E_{7,6})|_{\mathcal{A}}=x_4\partial_{x_7}+x_6\partial_{x_8}+x_9\partial_{x_{10}}+x_{11}\partial_{x_{13}}, \tag{2.49}$$

$$(E_{2,3}-E_{8,7})|_{\mathcal{A}}=x_3\partial_{x_4}+x_5\partial_{x_6}+x_{10}\partial_{x_{12}}+x_{13}\partial_{x_{14}}, \tag{2.50}$$

$$(E_{3,4}-E_{9,8})|_{\mathcal{A}}=-x_2\partial_{x_3}-x_6\partial_{x_9}-x_8\partial_{x_{10}}+x_{14}\partial_{x_{15}}, \tag{2.51}$$

$$(E_{4,5}-E_{10,9})|_{\mathcal{A}}=-x_1\partial_{x_2}+x_9\partial_{x_{11}}+x_{10}\partial_{x_{13}}+x_{12}\partial_{x_{14}}, \tag{2.52}$$

$$(E_{4,10}-E_{5,9})|_{\mathcal{A}}=-x_3\partial_{x_5}-x_4\partial_{x_6}-x_7\partial_{x_8}+x_{15}\partial_{x_{16}}, \tag{2.53}$$

$$(E_{1,3}-E_{8,6})|_{\mathcal{A}}=-x_3\partial_{x_7}-x_5\partial_{x_8}+x_9\partial_{x_{12}}+x_{11}\partial_{x_{14}}, \tag{2.54}$$

$$(E_{2,4}-E_{9,7})|_{\mathcal{A}}=x_2\partial_{x_4}-x_5\partial_{x_9}+x_8\partial_{x_{12}}+x_{13}\partial_{x_{15}}, \tag{2.55}$$

$$(E_{3,5}-E_{10,8})|_{\mathcal{A}}=-x_1\partial_{x_3}-x_6\partial_{x_{11}}-x_8\partial_{x_{13}}-x_{12}\partial_{x_{15}}, \tag{2.56}$$

$$(E_{3,10}-E_{5,8})|_{\mathcal{A}}=x_2\partial_{x_5}-x_4\partial_{x_9}-x_7\partial_{x_{10}}+x_{14}\partial_{x_{16}}, \tag{2.57}$$

$$(E_{1,4}-E_{9,6})|_{\mathcal{A}}=-x_2\partial_{x_7}+x_5\partial_{x_{10}}+x_6\partial_{x_{12}}+x_{11}\partial_{x_{15}}, \tag{2.58}$$

$$(E_{2,5}-E_{10,7})|_{\mathcal{A}}=x_1\partial_{x_4}-x_5\partial_{x_{11}}+x_8\partial_{x_{14}}-x_{10}\partial_{x_{15}}, \tag{2.59}$$

$$(E_{2,10}-E_{5,7})|_{\mathcal{A}}=-x_2\partial_{x_6}-x_3\partial_{x_9}+x_7\partial_{x_{12}}+x_{13}\partial_{x_{16}}, \tag{2.60}$$

$$(E_{3,9}-E_{4,8})|_{\mathcal{A}}=-x_1\partial_{x_5}+x_4\partial_{x_{11}}+x_7\partial_{x_{13}}+x_{12}\partial_{x_{16}}, \tag{2.61}$$

$$(E_{1,5}-E_{10,6})|_{\mathcal{A}}=-x_1\partial_{x_7}+x_5\partial_{x_{13}}+x_6\partial_{x_{14}}-x_9\partial_{x_{15}}, \tag{2.62}$$

$$(E_{1,10}-E_{5,6})|_{\mathcal{A}}=x_2\partial_{x_8}+x_3\partial_{x_{10}}+x_4\partial_{x_{12}}+x_{11}\partial_{x_{16}}, \tag{2.63}$$

$$(E_{2,9}-E_{4,7})|_{\mathcal{A}}=x_1\partial_{x_6}+x_3\partial_{x_{11}}-x_7\partial_{x_{14}}+x_{10}\partial_{x_{16}}, \tag{2.64}$$

$$(E_{1,9}-E_{4,6})|_{\mathcal{A}}=-x_1\partial_{x_8}-x_4\partial_{x_{14}}-x_3\partial_{x_{13}}+x_9\partial_{x_{16}}, \tag{2.65}$$

$$(E_{2,8}-E_{3,7})|_{\mathcal{A}}=x_1\partial_{x_9}-x_2\partial_{x_{11}}+x_7\partial_{x_{15}}-x_8\partial_{x_{16}}, \tag{2.66}$$

$$(E_{1,8}-E_{3,6})|_{\mathcal{A}}=-x_1\partial_{x_{10}}+x_2\partial_{x_{13}}+x_4\partial_{x_{15}}-x_6\partial_{x_{16}}, \tag{2.67}$$

$$(E_{1,7}-E_{2,6})|_{\mathcal{A}}=x_1\partial_{x_{12}}-x_2\partial_{x_{14}}+x_3\partial_{x_{15}}-x_5\partial_{x_{16}}, \tag{2.68}$$

$$(E_{j,i}-E_{5+i,5+j})|_{\mathcal{A}}=\tau[(E_{i,j}-E_{5+j,5+i})|_{\mathcal{A}}], \tag{2.69}$$

$$(E_{5+j,i}-E_{5+i,j})|_{\mathcal{A}}=\tau[(E_{i,5+j}-E_{j,5+i})|_{\mathcal{A}}] \tag{2.70}$$

for $1\leq i<j\leq 16$,

$$(E_{r,r}-E_{5+r,5+r})|_{\mathcal{A}}=\sum_{i=1}^{16}(1/2+a_{r,i})x_i\partial_{x_i},\qquad r\in\overline{1,5}, \tag{2.71}$$

where $a_{r,i}$ are given in the following table

Table 1

i	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$a_{1,i}$	0	0	0	0	0	0	-1	-1	0	-1	0	-1	-1	-1	-1	-1
$a_{2,i}$	0	0	0	-1	0	-1	0	0	-1	0	-1	-1	0	-1	-1	-1
$a_{3,i}$	0	0	-1	0	-1	0	0	0	-1	-1	-1	0	-1	0	-1	-1
$a_{4,i}$	0	-1	0	0	-1	-1	0	-1	0	0	-1	0	-1	-1	0	-1
$a_{5,i}$	-1	0	0	0	-1	-1	0	-1	-1	-1	0	-1	0	0	0	-1

Note that (2.39)-(2.68) are the representation formulas of all the positive root vectors. In particular, x_1 is a highest-weight vector of V with weight λ_4, the forth fundamental weight of E_6, and V gives a spin representation of $o(10,\mathbb{C})$.

3 Decomposition of the $o(10,\mathbb{C})$-Module $\mathcal{A}$

Recall that the representation of $\mathcal{G}^{D_5}$ on $\mathcal{A}$ is given by (2.37) and the representation of $o(10,\mathbb{C})$ is given in (2.48). We calculate

$$\alpha_r|_{\mathcal{A}} = \sum_{i=1}^{16} b_{r,i} x_i \partial_{x_i} \qquad \text{for } \ r \in \overline{1,5}, \tag{3.1}$$

where $b_{r,i}$ are given in the following table:

Table 2

i	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$b_{1,i}$	0	0	0	1	0	1	−1	−1	1	−1	1	0	−1	0	0	0
$b_{2,i}$	0	0	1	1	−1	−1	1	−1	0	0	0	0	0	0	1	−1
$b_{3,i}$	0	0	1	−1	1	−1	0	0	0	1	0	−1	1	−1	0	0
$b_{4,i}$	0	1	−1	0	0	1	0	1	−1	−1	0	0	0	1	−1	0
$b_{5,i}$	1	−1	0	0	0	0	0	0	1	1	−1	1	−1	−1	0	0

Recall that a singular vector of $o(10,\mathbb{C})$ is a nonzero weight vector annihilated by positive root vectors. Note that the weight of a singular vector in $\mathcal{A}$ must be dominate integral. The above table motivates us to assume that

$$\zeta_1 = a_1 x_1 x_{11} + a_2 x_2 x_9 + a_3 x_3 x_6 + a_4 x_4 x_5 \tag{3.2}$$

is a singular vector, where a_i are constants to be determined. By (2.49),

$$(E_{1,2} - E_{7,6})(\zeta_1) = 0. \tag{3.3}$$

Moreover, (2.50) says

$$(E_{2,3} - E_{8,7})(\zeta_1) = (a_3 + a_4) x_3 x_5 = 0 \Longrightarrow a_4 = -a_3. \tag{3.4}$$

Expression (2.51) gives

$$(E_{3,4} - E_{9,8})(\zeta_1) = -(a_2 + a_3) x_2 x_6 = 0 \Longrightarrow a_3 = -a_2. \tag{3.5}$$

Furthermore, (2.52) yields

$$(E_{4,5} - E_{10,9})(\zeta_1) = (a_1 - a_2) x_1 x_9 = 0 \Longrightarrow a_2 = a_1. \tag{3.6}$$

According to (2.53) and (3.4),

$$(E_{4,10} - E_{5,9})(\zeta_1) = -(a_3 + a_4) x_3 x_4 = 0. \tag{3.7}$$

Taking $a_1 = 1$, we have the singular vector

$$\zeta_1 = x_1 x_{11} + x_2 x_9 - x_3 x_6 + x_4 x_5 \tag{3.8}$$

of weight λ_1, the first fundamental weight of E_6. Thus ζ_1 generates the 10-dimensional natural $o(10,\mathbb{C})$-module U. According to (2.49)-(2.53), (2.69) and (2.70),

$$(E_{2,1} - E_{6,7})|_{\mathcal{A}} = x_7\partial_{x_4} + x_8\partial_{x_6} + x_{10}\partial_{x_9} + x_{13}\partial_{x_{11}}, \tag{3.9}$$

$$(E_{3,2} - E_{7,8})|_{\mathcal{A}} = x_4\partial_{x_3} + x_6\partial_{x_5} + x_{12}\partial_{x_{10}} + x_{14}\partial_{x_{13}}, \tag{3.10}$$

$$(E_{4,3} - E_{8,9})|_{\mathcal{A}} = -x_3\partial_{x_2} - x_9\partial_{x_6} - x_{10}\partial_{x_8} + x_{15}\partial_{x_{14}}, \tag{3.11}$$

$$(E_{5,4} - E_{9,10})|_{\mathcal{A}} = -x_2\partial_{x_1} + x_{11}\partial_{x_9} + x_{13}\partial_{x_{10}} + x_{14}\partial_{x_{12}}, \tag{3.12}$$

$$(E_{10,4} - E_{9,5})|_{\mathcal{A}} = -x_5\partial_{x_3} - x_6\partial_{x_4} - x_8\partial_{x_7} + x_{16}\partial_{x_{15}}. \tag{3.13}$$

We take

$$\zeta_2 = (E_{2,1} - E_{6,7})(\zeta_1) = x_1x_{13} + x_2x_{10} - x_3x_8 + x_5x_7, \tag{3.14}$$

$$\zeta_3 = (E_{3,2} - E_{7,8})(\zeta_2) = x_1x_{14} + x_2x_{12} - x_4x_8 + x_6x_7, \tag{3.15}$$

$$\zeta_4 = (E_{4,3} - E_{8,9})(\zeta_3) = x_1x_{15} - x_3x_{12} + x_4x_{10} - x_7x_9, \tag{3.16}$$

$$\zeta_5 = (E_{5,4} - E_{9,10})(\zeta_4) = -x_2x_{15} - x_3x_{14} + x_4x_{13} - x_7x_{11}, \tag{3.17}$$

$$\zeta_{10} = (E_{10,4} - E_{9,5})(\zeta_4) = x_1x_{16} + x_5x_{12} - x_6x_{10} + x_8x_9, \tag{3.18}$$

$$\zeta_9 = (E_{9,5} - E_{10,4})(\zeta_5) = x_2x_{16} - x_5x_{14} + x_6x_{13} - x_8x_{11}, \tag{3.19}$$

$$\zeta_8 = (E_{8,9} - E_{4,3})(\zeta_9) = x_3x_{16} + x_5x_{15} + x_9x_{13} - x_{10}x_{11}, \tag{3.20}$$

$$\zeta_7 = (E_{7,8} - E_{3,2})(\zeta_8) = -x_4x_{16} - x_6x_{15} - x_9x_{14} + x_{11}x_{12}, \tag{3.21}$$

$$\zeta_6 = (E_{6,7} - E_{2,1})(\zeta_7) = x_7x_{16} + x_8x_{15} + x_{10}x_{14} - x_{12}x_{13}. \tag{3.22}$$

Then $U = \sum_{i=1}^{10} \mathbb{C}\zeta_i$ forms an $o(10,\mathbb{C})$-module isomorphic to the 10-dimensional natural $o(10,\mathbb{C})$-module with $\{\zeta_1, ..., \zeta_{10}\}$ as the standard basis.

Theorem 3.1. *Any singular vector is a polynomial in x_1 and ζ_1.*

Proof. Note

$$x_{11} = x_1^{-1}(\zeta_1 - x_2x_9 + x_3x_6 - x_4x_5), \tag{3.23}$$

$$x_{13} = x_1^{-1}(\zeta_2 - x_2x_{10} + x_3x_8 - x_5x_7), \tag{3.24}$$

$$x_{14} = x_1^{-1}(\zeta_3 - x_2x_{12} + x_4x_8 - x_6x_7), \tag{3.25}$$

$$x_{15} = x_1^{-1}(\zeta_4 + x_3x_{12} - x_4x_{10} + x_7x_9), \tag{3.26}$$

$$x_{16} = x_1^{-1}(\zeta_{10} - x_5x_{12} + x_6x_{10} - x_8x_9). \tag{3.27}$$

Let f be a singular vector in $\mathcal{A}$. Substituting (3.23)-(3.27) into it, we can write

$$f = g(x_i, \zeta_1, \zeta_2, \zeta_3, \zeta_4, \zeta_{10} \mid 11, 13, 14, 15, 16 \neq i\overline{1,16}). \tag{3.28}$$

By (2.52), (2.56), (2.59), (2.61), (2.62) and (2.64)-(2.68),

$$(E_{4,5}-E_{10,9})(f)=-x_1\partial_{x_2}(g)=0\Longrightarrow g_{x_2}=0, \tag{3.29}$$

$$(E_{3,5}-E_{10,8})(f)=-x_1\partial_{x_3}(g)=0\Longrightarrow g_{x_3}=0, \tag{3.30}$$

$$(E_{2,5}-E_{10,7})(f)=x_1\partial_{x_4}(g)=0\Longrightarrow g_{x_4}=0, \tag{3.31}$$

$$(E_{3,9}-E_{4,8})(f)=-x_1\partial_{x_5}(g)=0\Longrightarrow g_{x_5}=0, \tag{3.32}$$

$$(E_{1,5}-E_{10,6})(f)=-x_1\partial_{x_7}(g)=0\Longrightarrow g_{x_7}=0, \tag{3.33}$$

$$(E_{2,9}-E_{4,7})(f)=x_1\partial_{x_6}(g)=0\Longrightarrow g_{x_6}=0, \tag{3.34}$$

$$(E_{1,9}-E_{4,6})(f)=-x_1\partial_{x_8}(g)=0\Longrightarrow g_{x_8}=0, \tag{3.35}$$

$$(E_{2,8}-E_{3,7})(f)=x_1\partial_{x_9}(g)=0\Longrightarrow g_{x_9}=0, \tag{3.36}$$

$$(E_{1,8}-E_{3,6})(f)=-x_1\partial_{x_{10}}(g)=0\Longrightarrow g_{x_{10}}=0, \tag{3.37}$$

$$(E_{1,7}-E_{2,6})(f)=x_1\partial_{x_{12}}(g)=0\Longrightarrow g_{x_{12}}=0. \tag{3.38}$$

Thus f is a function in $x_1,\zeta_1,\zeta_2,\zeta_3,\zeta_4$ and ζ_{10}.

According to (2.49)-(2.51) and (2.53),

$$(E_{1,2}-E_{7,6})(f)=\zeta_1\partial_{\zeta_2}(g)=0\Longrightarrow g_{\zeta_2}=0, \tag{3.39}$$

$$(E_{2,3}-E_{8,7})(f)=\zeta_2\partial_{\zeta_3}(g)=0\Longrightarrow g_{\zeta_3}=0, \tag{3.40}$$

$$(E_{3,4}-E_{9,8})(f)=\zeta_3\partial_{\zeta_4}(g)=0\Longrightarrow g_{\zeta_4}=0, \tag{3.41}$$

$$(E_{4,10}-E_{5,9})=\zeta_4\partial_{\zeta_{10}}(g)=0\Longrightarrow g_{\zeta_{10}}=0. \tag{3.42}$$

Hence f is a function in x_1 and ζ_1. Thanks to (3.23), it must be a polynomial in x_1 and ζ_1. $\square$

Let V_{m_1,m_2} be the irreducible $o(10,\mathbb{C})$-submodule generated by $x_1^{m_1}\zeta^{m_2}$. By Weyl's theorem of complete reducibility,

$$\mathcal{A}=\bigoplus_{m_1,m_2=0}^{\infty}V_{m_1,m_2}. \tag{3.43}$$

Denote by $V(\lambda)$ the highest-weight irreducible $o(10,\mathbb{C})$-module with the highest weight λ. The above equation leads to the following combinatorial identity:

$$\sum_{m_1,m_2=0}^{\infty}(\dim V(m_2\lambda_1+m_1\lambda_4))q^{m_1+2m_2}=\frac{1}{(1-q)^{16}}, \tag{3.44}$$

which was proved in (3.23)-(3.41) by partial differential equations.

4 Realization of E_6 in 16-Dimensional Space

In this section, we want to find a differential-operator representation of $\mathcal{G}^{E_6}$, equivalently, a fraction representation on 16-dimensional space of the Lie group of type E_6.

According to (2.36) and (2.37), we calculate

$$\alpha_6|_{\mathcal{A}} = -2x_1\partial_{x_1} - \sum_{i=2}^{10} x_i\partial_{x_i} - x_{12}\partial_{x_{12}}. \tag{4.1}$$

Write

$$\widehat{\alpha} = 2\alpha_1 + 4\alpha_3 + 6\alpha_4 + 3\alpha_2 + 5\alpha_5 + 4\alpha_6. \tag{4.2}$$

Then

$$(\widehat{\alpha}, \alpha_r) = 0 \qquad \text{for } \ r \in \overline{1,5} \tag{4.3}$$

by the Dynkin diagram of E_6. Thanks to (2.9),

$$[\widehat{\alpha}, \mathcal{G}^{D_5}] = 0. \tag{4.4}$$

By Schur's Lemma, $\widehat{\alpha}|_V = c\sum_{i=1}^{16} x_i\partial_{x_i}$. According to the coefficients of $x_1\partial_{x_1}$ in (3.1) with the data in Table 2 and (4.2), we have

$$\widehat{\alpha}|_{\mathcal{A}} = \sum_{i=1}^{16} x_i\partial_{x_i} = D, \tag{4.5}$$

the degree operator on $\mathcal{A}$.

Recall that the Lie bracket in the algebra $\mathbb{A}$ (cf. (2.40)) is given by the commutator

$$[d_1, d_2] = d_1d_2 - d_2d_1. \tag{4.6}$$

Set

$$\mathcal{D} = \sum_{i=1}^{16} \mathbb{C}\partial_{x_i}. \tag{4.7}$$

Then $\mathcal{D}$ forms an $o(10,\mathbb{C})$-module with respect to the action

$$B(d) = [B|_{\mathcal{A}}, \partial] \qquad \text{for } \ B \in o(10,\mathbb{C}),\ \partial \in \mathcal{D}. \tag{4.8}$$

On the other hand, $\mathcal{G}_{\pm}$ (cf. (2.22)-(2.32)) form $o(10,\mathbb{C})$-modules with respect to the action

$$B(u) = [\nu(B), u] \qquad \text{for } \ B \in o(10,\mathbb{C}),\ u \in \mathcal{G}_{\pm}. \tag{4.9}$$

According to (2.36) and (2.37), the linear map determined by $\eta_i \mapsto x_i$ for $i \in \overline{1,16}$ gives an $o(10,\mathbb{C})$-module isomorphism from $\mathcal{G}_-$ to V. Moreover, (2.12) and (2.34) implies that the linear map determined by $\xi_i \mapsto \partial_{x_i}$ for $i \in \overline{1,16}$ gives an $o(10,\mathbb{C})$-module isomorphism from $\mathcal{G}_+$ to $\mathcal{D}$. Hence we define the action of $\mathcal{G}_+$ on $\mathcal{A}$ by

$$\xi_i|_{\mathcal{A}} = \partial_{x_i} \qquad \text{for } \ i \in \overline{1,16}. \tag{4.10}$$

Recall the Witt Lie subalgebra of $\mathbb{A}$:

$$\mathcal{W}_{16} = \sum_{i=1}^{16} \mathcal{A}\partial_{x_i}. \tag{4.11}$$

Now we want to find the differential operators $P_1, P_2, ..., P_{16} \in \mathcal{W}_{16}$ such that the following action matches the structure of $\mathcal{G}^{E_6}$:

$$\eta_i|_{\mathcal{A}} = P_i \qquad \text{for } \ i \in \overline{1,16}. \tag{4.12}$$

Imposing

$$[\partial_{x_1}, P_1] = [E_{\alpha_6}, E_{-\alpha_6}]|_{\mathcal{A}} = -\alpha_6|_{\mathcal{A}} = 2x_1\partial_{x_1} + \sum_{i=2}^{10} x_i\partial_{x_i} + x_{12}\partial_{x_{12}}, \tag{4.13}$$

we take

$$P_1 = x_1(\sum_{i=1}^{10} x_i\partial_{x_i} + x_{12}\partial_{x_{12}}) + P_1', \tag{4.14}$$

where P_1' is a differential operator such that $[\partial_{x_1}, P_1'] = 0$. Moreover,

$$[\partial_{x_r}, x_1(\sum_{i=1}^{10} x_i\partial_{x_i} + x_{12}\partial_{x_{12}})] = x_1\partial_{x_r} \qquad \text{for } \ r \in \{\overline{2,10}, 12\}. \tag{4.15}$$

Wanting $[\partial_{x_r}, P_1] \in \mathcal{G}^{D_5}|_{\mathcal{A}} = o(10,\mathbb{C})|_{\mathcal{A}}$ (cf. (2.49)-(2.71)), we take

$$\begin{aligned} P_1 &= x_1(\sum_{i=1}^{10} x_i\partial_{x_i} + x_{12}\partial_{x_{12}}) - (x_2x_9 - x_3x_6 + x_4x_5)\partial_{x_{11}} \\ &\quad -(x_2x_{10} - x_3x_8 + x_5x_7)\partial_{x_{13}} - (x_2x_{12} - x_4x_8 + x_6x_7)\partial_{x_{14}} \\ &\quad +(x_3x_{12} - x_4x_{10} + x_7x_9)\partial_{x_{15}} - (x_5x_{12} - x_6x_{10} + x_8x_9)\partial_{x_{16}} \\ &= x_1D - \zeta_1\partial_{x_{11}} - \zeta_2\partial_{x_{13}} - \zeta_3\partial_{x_{14}} - \zeta_4\partial_{x_{15}} - \zeta_{10}\partial_{x_{16}} \end{aligned} \tag{4.16}$$

by (2.52), (2.56), (2.59), (2.61), (2.62), (2.64)-(2.68), (3.8), (3.14)-(3.16), (3.18) and (4.5). Then

$$[\partial_{x_s}, P_1] = [\xi_s, \eta_1]|_{\mathcal{A}} \qquad \text{for } \ s \in \overline{1,16} \tag{4.17}$$

due to (2.42)-(2.48).

Since $[E_{-\alpha_5}, \eta_1] = \eta_2$ by (2.10), we take

$$\begin{aligned} P_2 &= [E_{-\alpha_5}|_{\mathcal{A}}, \eta_1|_{\mathcal{A}}] = -[(E_{5,4} - E_{9,10})|_{\mathcal{A}}, P_1] \\ &= x_2D - \zeta_1\partial_{x_9} - \zeta_2\partial_{x_{10}} - \zeta_3\partial_{x_{12}} + \zeta_5\partial_{x_{15}} - \zeta_9\partial_{x_{16}} \end{aligned} \tag{4.18}$$

by (2.44) and (3.12). Note that $[E_{-\alpha_4},\eta_2]=\eta_3$ by (2.10). Hence (3.11) gives

$$\begin{aligned} P_3 &= [E_{-\alpha_4}|_{\mathcal{A}},\eta_2|_{\mathcal{A}}]=-[(E_{4,3}-E_{8,9})|_{\mathcal{A}},P_2] \\ &= x_3D+\zeta_1\partial_{x_6}+\zeta_2\partial_{x_8}+\zeta_4\partial_{x_{12}}+\zeta_5\partial_{x_{14}}-\zeta_8\partial_{x_{16}}. \end{aligned}\tag{4.19}$$

Thanks to $-[E_{-\alpha_3},\eta_3]=\eta_4$, (2.50) and (3.10), we have

$$\begin{aligned} P_4 &= -[E_{-\alpha_3}|_{\mathcal{A}},\eta_3|_{\mathcal{A}}]=[(E_{3,2}-E_{7,9})|_{\mathcal{A}},P_3] \\ &= x_4D-\zeta_1\partial_{x_5}+\zeta_3\partial_{x_8}-\zeta_4\partial_{x_{10}}-\zeta_5\partial_{x_{13}}+\zeta_7\partial_{x_{16}}. \end{aligned}\tag{4.20}$$

Observe that $[E_{-\alpha_2},\eta_3]=\eta_5$ by (2.10). So (3.13) yields

$$\begin{aligned} P_5 &= [E_{-\alpha_2}|_{\mathcal{A}},\eta_3|_{\mathcal{A}}]=-[(E_{10,4}-E_{9,5})|_{\mathcal{A}},P_2] \\ &= x_5D-\zeta_1\partial_{x_4}-\zeta_2\partial_{x_7}-\zeta_{10}\partial_{x_{12}}+\zeta_9\partial_{x_{14}}-\zeta_8\partial_{x_{15}}. \end{aligned}\tag{4.21}$$

Since $[E_{-\alpha_2},\eta_4]=\eta_6$, (2.13) implies

$$\begin{aligned} P_6 &= [E_{-\alpha_2}|_{\mathcal{A}},\eta_4|_{\mathcal{A}}]=-[(E_{10,4}-E_{9,5})|_{\mathcal{A}},P_4] \\ &= x_6D+\zeta_1\partial_{x_3}-\zeta_3\partial_{x_7}+\zeta_{10}\partial_{x_{10}}-\zeta_9\partial_{x_{13}}+\zeta_7\partial_{x_{15}}. \end{aligned}\tag{4.22}$$

As $-[E_{-\alpha_1},\eta_4]=\eta_7$, we get by (3.9) that

$$\begin{aligned} P_7 &= -[E_{-\alpha_1}|_{\mathcal{A}},\eta_4|_{\mathcal{A}}]=[(E_{2,1}-E_{6,7})|_{\mathcal{A}},P_4] \\ &= x_7D-\zeta_2\partial_{x_5}-\zeta_3\partial_{x_6}+\zeta_4\partial_{x_9}+\zeta_5\partial_{x_{11}}-\zeta_6\partial_{x_{16}}. \end{aligned}\tag{4.23}$$

Thanks to $[E_{-\alpha_2},\eta_7]=\eta_8$, (3.13) gives

$$\begin{aligned} P_8 &= [E_{-\alpha_2}|_{\mathcal{A}},\eta_7|_{\mathcal{A}}=-[(E_{10,4}-E_{9,5})|_{\mathcal{A}},P_7] \\ &= x_8D+\zeta_2\partial_{x_3}+\zeta_3\partial_{x_4}-\zeta_{10}\partial_{x_9}+\zeta_9\partial_{x_{11}}-\zeta_6\partial_{x_{15}}. \end{aligned}\tag{4.24}$$

The fact $[E_{-\alpha_4},\eta_6]=\eta_9$ yields

$$\begin{aligned} P_9 &= [E_{-\alpha_4}|_{\mathcal{A}},\eta_6|_{\mathcal{A}}]=-[(E_{4,3}-E_{8,9})|_{\mathcal{A}},P_6] \\ &= x_9D-\zeta_1\partial_{x_2}+\zeta_4\partial_{x_7}-\zeta_{10}\partial_{x_8}-\zeta_8\partial_{x_{13}}+\zeta_7\partial_{x_{14}}. \end{aligned}\tag{4.25}$$

by (3.11). As $[E_{-\alpha_4},\eta_8]=\eta_{10}$, we find

$$\begin{aligned} P_{10} &= [E_{-\alpha_4}|_{\mathcal{A}},\eta_8|_{\mathcal{A}}=-[(E_{4,3}-E_{8,9})|_{\mathcal{A}},P_8] \\ &= x_{10}D-\zeta_2\partial_{x_2}-\zeta_4\partial_{x_4}+\zeta_{10}\partial_{x_6}+\zeta_8\partial_{x_{11}}-\zeta_6\partial_{x_{14}}. \end{aligned}\tag{4.26}$$

by (3.11). Moreover, the fact $-[E_{-\alpha_5},\eta_9]=\eta_{11}$ implies

$$\begin{aligned} P_{11} &= -[E_{-\alpha_5}|_{\mathcal{A}},\eta_9|_{\mathcal{A}}]=[(E_{5,4}-E_{9,10})|_{\mathcal{A}},P_9] \\ &= x_{11}D-\zeta_1\partial_{x_1}+\zeta_5\partial_{x_7}+\zeta_9\partial_{x_8}+\zeta_8\partial_{x_{10}}-\zeta_7\partial_{x_{12}}. \end{aligned}\tag{4.27}$$

by (3.12). Since $-[E_{-\alpha_3},\eta_{10}]=\eta_{12}$,

$$\begin{aligned} P_{12} &= -[E_{-\alpha_3}|_{\mathcal{A}},\eta_{10}|_{\mathcal{A}}=[(E_{3,2}-E_{7,8})|_{\mathcal{A}},P_{10}] \\ &= x_{12}D-\zeta_3\partial_{x_2}+\zeta_4\partial_{x_3}-\zeta_{10}\partial_{x_5}-\zeta_7\partial_{x_{11}}+\zeta_6\partial_{x_{13}} \end{aligned}\tag{4.28}$$

by (3.10).

Observing $-[E_{-\alpha_1},\eta_{11}]=\eta_{13}$, we have

$$\begin{aligned} P_{13} &= -[E_{-\alpha_1}|_{\mathcal{A}},\eta_{11}|_{\mathcal{A}}]=[(E_{2,1}-E_{6,7})|_{\mathcal{A}},P_{11}] \\ &= x_{13}D-\zeta_2\partial_{x_1}-\zeta_5\partial_{x_4}-\zeta_9\partial_{x_6}-\zeta_8\partial_{x_9}+\zeta_6\partial_{x_{12}}. \end{aligned}\tag{4.29}$$

by (3.9). The fact $-[E_{-\alpha_3},\eta_{13}]=\eta_{14}$ gives

$$\begin{aligned} P_{14} &= -[E_{-\alpha_3}|_{\mathcal{A}},\eta_{13}|_{\mathcal{A}}]=[(E_{3,2}-E_{7,8})|_{\mathcal{A}},P_{13}] \\ &= x_{14}D-\zeta_3\partial_{x_1}+\zeta_5\partial_{x_3}+\zeta_9\partial_{x_5}+\zeta_7\partial_{x_9}-\zeta_6\partial_{x_{10}}. \end{aligned}\tag{4.30}$$

by (3.10). As $-[E_{-\alpha_4},\eta_{14}]=\eta_{15}$, we get

$$\begin{aligned} P_{15} &= -[E_{-\alpha_4}|_{\mathcal{A}},\eta_{14}|_{\mathcal{A}}]=[(E_{4,3}-E_{8,9})|_{\mathcal{A}},P_{14}] \\ &= x_{15}D-\zeta_4\partial_{x_1}+\zeta_5\partial_{x_2}-\zeta_8\partial_{x_5}+\zeta_7\partial_{x_6}-\zeta_6\partial_{x_8}. \end{aligned}\tag{4.31}$$

by (3.11). Since $-[E_{-\alpha_2},\eta_{15}]=\eta_{16}$, we have

$$\begin{aligned} P_{16} &= -[E_{-\alpha_2}|_{\mathcal{A}},\eta_{15}|_{\mathcal{A}}]=[(E_{10,4}-E_{9,5})|_{\mathcal{A}},P_{15}] \\ &= x_{16}D-\zeta_{10}\partial_{x_1}-\zeta_9\partial_{x_2}-\zeta_8\partial_{x_3}+\zeta_7\partial_{x_4}-\zeta_6\partial_{x_7}. \end{aligned}\tag{4.32}$$

by (3.13).

Set

$$\mathcal{P}=\sum_{i=1}^{16}\mathbb{C}P_i,\qquad \mathcal{C}_0=o(10,\mathbb{C})|_{\mathcal{A}}+\mathbb{C}D\tag{4.33}$$

(cf. (2.49)-(2.71) and (4.5)) and

$$\mathcal{C}=\mathcal{P}+\mathcal{C}_0+\mathcal{D}\tag{4.34}$$

(cf. (4.7)). The we have:

Theorem 4.1. *The space $\mathcal{C}$ forms a Lie subalgebra of the Witt algebra $\mathcal{W}_{16}$ (cf. (4.11)). Moreover, the linear map ϑ determined by*

$$\vartheta(\xi_i)=\partial_{x_i},\;\;\vartheta(\eta_i)=P_i,\;\;\vartheta(u)=\nu^{-1}(u)|_{\mathcal{A}}\qquad \textit{for}\;\; i\in\overline{1,16},\;u\in\mathcal{G}^{D_5}\tag{4.35}$$

(cf. (2.42)-(2.47)) and

$$\vartheta(\alpha_6)=-2x_1\partial_{x_1}-\sum_{i=2}^{10}x_i\partial_{x_i}-x_{12}\partial_{x_{12}}\tag{4.36}$$

(cf. (4.1)) gives a Lie algebra isomorphism from $\mathcal{G}^{E_6}$ to $\mathcal{C}$.

Proof. Since $\mathcal{D}\cong\mathcal{G}_+$ as $\mathcal{G}^{D_5}$-modules, we have

$$\mathcal{G}_0+\mathcal{G}_+\overset{\vartheta}{\cong}\mathcal{C}_0+\mathcal{D} \tag{4.37}$$

as Lie algebras. Denote by $U(\mathcal{G})$ the universal enveloping algebra of a Lie algebra $\mathcal{G}$. Note that

$$\mathcal{B}_-=\mathcal{G}_0+\mathcal{G}_-,\qquad \mathcal{B}_+=\mathcal{G}_0+\mathcal{G}_+ \tag{4.38}$$

are also Lie subalgebras of $\mathcal{G}^{E_6}$ and

$$\mathcal{G}^{E_6}=\mathcal{B}_-\oplus\mathcal{G}_+=\mathcal{G}_-\oplus\mathcal{B}_+. \tag{4.39}$$

We define a one-dimensional $\mathcal{B}_-$-module $\mathbb{C}u_0$ by

$$w(u_0)=0\qquad\text{for }\; w\in\mathcal{B}_-\bigcap\mathcal{G}^{D_5},\;\; \widehat{\alpha}(u_0)=-16u_0 \tag{4.40}$$

(cf. (4.2)). Let

$$\Psi=U(\mathcal{G}^{E_6})\otimes_{\mathcal{B}_-}\mathbb{C}u_0\cong U(\mathcal{G}_+)\otimes_{\mathbb{C}}\mathbb{C}u_0 \tag{4.41}$$

be the induced $\mathcal{G}^{E_6}$-module.

Reacll that $\mathbb{N}$ is the set of nonnegative integers. Let

$$\mathcal{A}'=\mathbb{C}[\partial_{x_1},\partial_{x_2},...,\partial_{x_{16}}]. \tag{4.42}$$

We define an action of the associative algebra $\mathbb{A}$ (cf. (2.40)) on $\mathcal{A}'$ by

$$\partial_{x_i}(\prod_{j=1}^{16}\partial_{x_j}^{\beta_j})=\partial_{x_i}^{\beta_i+1}\prod_{i\neq j\in\overline{1,16}}\prod_{j=1}^{16}\partial_{x_j}^{\beta_j} \tag{4.43}$$

and

$$x_i(\prod_{j=1}^{16}\partial_{x_j}^{\beta_j})=-\beta_i\partial_{x_i}^{\beta_i-1}\prod_{i\neq j\in\overline{1,16}}\prod_{j=1}^{16}\partial_{x_j}^{\beta_j} \tag{4.44}$$

for $i\in\overline{1,16}$. Since

$$[-x_i,\partial_{x_j}]=[\partial_{x_i},x_j]=\delta_{i,j}\qquad\text{for }\; i,j\in\overline{1,16}, \tag{4.45}$$

the above action gives an associative algebra representation of $\mathbb{A}$. Thus it also gives a Lie algebra representation of $\mathbb{A}$ (cf. (4.6)). It is straightforward to verify that

$$[d|_{\mathcal{A}'},\partial|_{\mathcal{A}'}]=[d,\partial]|_{\mathcal{A}'}\qquad\text{for }\; d\in\mathcal{C}_0,\;\partial\in\mathcal{D}. \tag{4.46}$$

Define linear map $\varsigma:\Psi\rightarrow\mathcal{A}'$ by

$$\varsigma(\prod_{i=1}^{16}\xi_i^{\ell_i}\otimes u_0)=\prod_{i=1}^{16}\partial_{x_i}^{\ell_i}\qquad(\ell_1,...,\ell_{16})\in\mathbb{N}^{16}. \tag{4.47}$$

According to (2.49)-(2.71), (4.43) and (4.44),

$$D(1)=-16,\;\; d(1)=0 \qquad \text{for}\;\; d\in o(10,\mathbb{C})|_{\mathcal{A}}. \tag{4.48}$$

Moreover, (4.40), (4.41), (4.43), (4.44) and (4.48) imply

$$\varsigma(\xi(v))=\vartheta(\xi)\varsigma(v) \qquad \text{for}\;\; \xi\in\mathcal{G}_0,\;\; v\in\Psi. \tag{4.49}$$

Now (4.41) and (4.43) imply

$$\varsigma(w(u))=\vartheta(w)(\varsigma(u)) \qquad \text{for}\;\; w\in\mathcal{B}_+,\;\;\; u\in\Psi. \tag{4.50}$$

Thus we have

$$\varsigma w|_{\Psi}\varsigma^{-1}=\vartheta(w)|_{\mathcal{A}'} \qquad \text{for}\;\; w\in\mathcal{B}_+. \tag{4.51}$$

On the other hand, the linear map

$$\psi(v)=\varsigma v|_{\Psi}\varsigma^{-1} \qquad \text{for}\;\; v\in\mathcal{G}^{E_6} \tag{4.52}$$

is a Lie algebra monomorphism from $\mathcal{G}^{E_6}$ to $\mathbb{A}|_{\mathcal{A}'}$. According to (4.17) and (4.45),

$$\psi(\eta_1)=P_1|_{\mathcal{A}'}. \tag{4.53}$$

By the constructions of $P_2,...,P_{16}$ in (4.18)-(4.32), we have

$$\psi(\eta_i)=P_i|_{\mathcal{A}'} \qquad \text{for}\;\; i\in\overline{2,16}. \tag{4.54}$$

Therefore, we have

$$\psi(v)=\vartheta(v)|_{\mathcal{A}'} \qquad \text{for}\;\; v\in\mathcal{G}^{E_6}. \tag{4.55}$$

In particular, $\mathcal{C}|_{\mathcal{A}'}=\vartheta(\mathcal{G}^{E_6})|_{\mathcal{A}'}=\psi(\mathcal{G}^{E_6})$ forms a Lie algebra. Since the linear map $d\mapsto d|_{\mathcal{A}'}$ for $d\in\mathcal{C}$ is injective, we have that $\mathcal{C}$ forms a Lie subalgebra of $\mathbb{A}$ and ϑ is a Lie algebra isomorphism. $\qquad\Box$

By the above theorem, a Lie group of type E_6 is generated by the real spinor transformations corresponding to (2.49)-(2.71), the real translations and dilations in $\sum_{i=1}^{16}\mathbb{R}x_i$, and the following fractional transformations:

$$\wp_{1b}(x_i)=\frac{x_i}{1-bx_1},\;\;\; i\in\{\overline{1,10},12\},\;\;\; \wp_{1b}(x_{11})=x_{11}-\frac{b(x_2x_9-x_3x_6+x_4x_5)}{1-bx_1}, \tag{4.56}$$

$$\wp_{1b}(x_{13})=x_{13}-\frac{b(x_2x_{10}-x_3x_8+x_5x_7)}{1-bx_1},\;\; \wp_{1b}(x_{14})=x_{14}-\frac{b(x_2x_{12}-x_4x_8+x_6x_7)}{1-bx_1}, \tag{4.57}$$

$$\wp_{1b}(x_{15})=x_{15}+\frac{b(x_3x_{12}-x_4x_{10}+x_7x_9)}{1-bx_1}, \tag{4.58}$$

$$\wp_{1b}(x_{16}) = x_{16} - \frac{b(x_5x_{12} - x_6x_{10} + x_8x_9)}{1 - bx_1}; \tag{4.59}$$

$$\wp_{2b}(x_i) = \frac{x_i}{1 - bx_2}, \quad i \in \overline{1,14} \setminus \{9, 10, 12\}, \quad \wp_{2b}(x_9) = x_9 - \frac{b(x_1x_{11} - x_3x_6 + x_4x_5)}{1 - bx_2}, \tag{4.60}$$

$$\wp_{2b}(x_{10}) = x_{10} - \frac{b(x_1x_{13} - x_3x_8 + x_5x_7)}{1 - bx_2}, \ \wp_{2b}(x_{12}) = x_{12} - \frac{b(x_1x_{14} - x_4x_8 + x_6x_7)}{1 - bx_2}, \tag{4.61}$$

$$\wp_{2b}(x_{15}) = x_{15} - \frac{b(x_3x_{14} - x_4x_{13} + x_7x_{11})}{1 - bx_2}, \tag{4.62}$$

$$\wp_{2b}(x_{16}) = x_{16} + \frac{b(x_5x_{14} - x_6x_{13} + x_8x_{11})}{1 - bx_2}; \tag{4.63}$$

$$\wp_{3b}(x_i) = \frac{x_i}{1 - bx_3}, \ i \in \overline{1,15} \setminus \{6, 8, 12, 14\}, \ \wp_{3b}(x_6) = x_6 + \frac{b(x_1x_{11} + x_2x_9 + x_4x_5)}{1 - bx_3}, \tag{4.64}$$

$$\wp_{3b}(x_8) = x_8 + \frac{b(x_1x_{13} + x_2x_{10} + x_5x_7)}{1 - bx_3}, \ \wp_{3b}(x_{12}) = x_{12} + \frac{b(x_1x_{15} + x_4x_{10} - x_7x_9)}{1 - bx_3}, \tag{4.65}$$

$$\wp_{3b}(x_{14}) = x_{14} - \frac{b(x_2x_{15} - x_4x_{13} + x_7x_{11})}{1 - bx_3}, \tag{4.66}$$

$$\wp_{3b}(x_{16}) = x_{16} - \frac{b(x_5x_{15} + x_9x_{13} - x_{10}x_{11})}{1 - bx_3}; \tag{4.67}$$

$$\wp_{4b}(x_i) = \frac{x_i}{1 - bx_4}, \ i \in \overline{1,15} \setminus \{5, 8, 10, 13\}, \ \wp_{4b}(x_5) = x_5 - \frac{b(x_1x_{11} + x_2x_9 - x_3x_6)}{1 - bx_4}, \tag{4.68}$$

$$\wp_{4b}(x_8) = x_8 + \frac{b(x_1x_{14} + x_2x_{12} + x_6x_7)}{1 - bx_4}, \ \wp_{4b}(x_{10}) = x_{10} - \frac{b(x_1x_{15} - x_3x_{12} - x_7x_9)}{1 - bx_4}, \tag{4.69}$$

$$\wp_{4b}(x_{13}) = x_{13} + \frac{b(x_2x_{15} + x_3x_{14} + x_7x_{11})}{1 - bx_4}, \tag{4.70}$$

$$\wp_{4b}(x_{16}) = x_{16} - \frac{b(x_6x_{15} + x_9x_{14} - x_{11}x_{12})}{1 - bx_4}; \tag{4.71}$$

$$\wp_{5b}(x_i) = \frac{x_i}{1 - bx_5}, \quad \wp_{5b}(x_4) = x_4 - \frac{b(x_1x_{11} + x_2x_9 - x_3x_6)}{1 - bx_5}, \tag{4.72}$$

$i \in \overline{1,16} \setminus \{4, 7, 12, 14, 15\}$,

$$\wp_{5b}(x_7) = x_7 - \frac{b(x_1x_{13} + x_2x_{10} - x_3x_8)}{1 - bx_5}, \ \wp_{5b}(x_{12}) = x_{12} - \frac{b(x_1x_{16} - x_6x_{10} + x_8x_9)}{1 - bx_5}, \tag{4.73}$$

$$\wp_{5b}(x_{14}) = x_{14} + \frac{b(x_2x_{16} + x_6x_{13} - x_8x_{11})}{1 - bx_5}, \tag{4.74}$$

$$\wp_{5b}(x_{15}) = x_{15} - \frac{b(x_3x_{16} + x_9x_{13} - x_{10}x_{11})}{1 - bx_5}; \tag{4.75}$$

$$\wp_{6b}(x_i) = \frac{x_i}{1 - bx_6}, \quad \wp_{6b}(x_3) = x_3 + \frac{b(x_1x_{11} + x_2x_9 + x_4x_5)}{1 - bx_6}, \tag{4.76}$$

$i \in \overline{1,16} \setminus \{3, 7, 10, 13, 15\}$,

$$\wp_{6b}(x_7) = x_7 - \frac{b(x_1x_{14} + x_2x_{12} - x_4x_8)}{1 - bx_6}, \ \wp_{6b}(x_{10}) = x_{10} + \frac{b(x_1x_{16} + x_5x_{12} + x_8x_9)}{1 - bx_6}, \tag{4.77}$$

$$\wp_{6b}(x_{13}) = x_{13} - \frac{b(x_2x_{16} - x_5x_{14} - x_8x_{11})}{1 - bx_6}, \tag{4.78}$$

$$\wp_{6b}(x_{15}) = x_{15} - \frac{b(x_4x_{16} + x_9x_{14} - x_{11}x_{12})}{1 - bx_6}; \tag{4.79}$$

$$\wp_{7b}(x_i) = \frac{x_i}{1 - bx_7},\ i \in \overline{1,15} \backslash \{5,6,9,11\},\ \wp_{7b}(x_5) = x_5 - \frac{b(x_1x_{13} + x_2x_{10} - x_3x_8)}{1 - bx_7}, \tag{4.80}$$

$$\wp_{7b}(x_6) = x_6 - \frac{b(x_1x_{14} + x_2x_{12} - x_4x_8)}{1 - bx_7},\ \wp_{7b}(x_9) = x_9 + \frac{b(x_1x_{15} - x_3x_{12} + x_4x_{10})}{1 - bx_7}, \tag{4.81}$$

$$\wp_{7b}(x_{11}) = x_{11} - \frac{b(x_2x_{15} + x_3x_{14} - x_4x_{13})}{1 - bx_7}, \tag{4.82}$$

$$\wp_{7b}(x_{16}) = x_{16} - \frac{b(x_8x_{15} + x_{10}x_{14} - x_{12}x_{13})}{1 - bx_7}; \tag{4.83}$$

$$\wp_{8b}(x_i) = \frac{x_i}{1 - bx_8},\quad \wp_{8b}(x_3) = x_3 + \frac{b(x_1x_{13} - x_3x_8 + x_5x_7)}{1 - bx_8}, \tag{4.84}$$

$i \in \overline{1,16} \setminus \{3,4,9,11,15\}$,

$$\wp_{8b}(x_4) = x_4 + \frac{b(x_1x_{14} + x_2x_{12} + x_6x_7)}{1 - bx_8},\ \wp_{8b}(x_9) = x_9 - \frac{b(x_1x_{16} + x_5x_{12} - x_6x_{10})}{1 - bx_8}, \tag{4.85}$$

$$\wp_{8b}(x_{11}) = x_{11} + \frac{b(x_2x_{16} - x_5x_{14} + x_6x_{13})}{1 - bx_8}, \tag{4.86}$$

$$\wp_{8b}(x_{15}) = x_{15} - \frac{b(x_7x_{16} + x_{10}x_{14} - x_{12}x_{13})}{1 - bx_8}; \tag{4.87}$$

$$\wp_{9b}(x_i) = \frac{x_i}{1 - bx_9},\quad \wp_{9b}(x_2) = x_2 - \frac{b(x_1x_{11} - x_3x_6 + x_4x_5)}{1 - bx_9}, \tag{4.88}$$

$i \in \overline{1,16} \setminus \{2,7,8,13,14\}$,

$$\wp_{9b}(x_7) = x_7 + \frac{b(x_1x_{15} - x_3x_{12} + x_4x_{10})}{1 - bx_9},\ \wp_{9b}(x_8) = x_8 - \frac{b(x_1x_{16} + x_5x_{12} - x_6x_{10})}{1 - bx_9}, \tag{4.89}$$

$$\wp_{9b}(x_{13}) = x_{13} - \frac{b(x_3x_{16} + x_5x_{15} - x_{10}x_{11})}{1 - bx_9}, \tag{4.90}$$

$$\wp_{9b}(x_{14}) = x_{14} - \frac{b(x_4x_{16} + x_6x_{15} - x_{11}x_{12})}{1 - bx_9}; \tag{4.91}$$

$$\wp_{10b}(x_i) = \frac{x_i}{1 - bx_{10}},\quad \wp_{10b}(x_2) = x_2 - \frac{b(x_1x_{13} - x_3x_8 + x_5x_7)}{1 - bx_{10}}, \tag{4.92}$$

$i \in \overline{1,16} \setminus \{2,4,6,11,14\}$,

$$\wp_{10b}(x_4) = x_4 - \frac{b(x_1x_{15} - x_3x_{12} - x_7x_9)}{1 - bx_{10}}, \tag{4.93}$$

$$\wp_{10b}(x_6) = x_6 + \frac{b(x_1x_{16} + x_5x_{12} + x_8x_9)}{1 - bx_{10}}, \tag{4.94}$$

$$\wp_{10b}(x_{11}) = x_{11} + \frac{b(x_3x_{16} + x_5x_{15} + x_9x_{13})}{1 - bx_{11}}, \tag{4.95}$$

$$\wp_{10b}(x_{14}) = x_{14} - \frac{b(x_7x_{16} + x_8x_{15} - x_{12}x_{13})}{1 - bx_{10}}; \tag{4.96}$$

$$\wp_{11b}(x_i) = \frac{x_i}{1 - bx_{11}}, \quad \wp_{11b}(x_1) = x_1 - \frac{b(x_2x_9 - x_3x_6 + x_4x_5)}{1 - bx_{11}}, \tag{4.97}$$

$i \in \overline{1,16} \setminus \{1,7,8,10,12\}$,

$$\wp_{11b}(x_7) = x_7 - \frac{b(x_2x_{15} + x_3x_{14} - x_4x_{13})}{1 - bx_{11}}, \tag{4.98}$$

$$\wp_{11b}(x_8) = x_8 + \frac{b(x_2x_{16} - x_5x_{14} + x_6x_{13})}{1 - bx_{11}}, \tag{4.99}$$

$$\wp_{11b}(x_{10}) = x_{10} + \frac{b(x_3x_{16} + x_5x_{15} + x_9x_{13})}{1 - bx_{11}}, \tag{4.100}$$

$$\wp_{11b}(x_{12}) = x_{12} + \frac{b(x_4x_{16} + x_6x_{15} + x_9x_{14})}{1 - bx_{11}}; \tag{4.101}$$

$$\wp_{12b}(x_i) = \frac{x_i}{1 - bx_{12}}, \quad \wp_{12b}(x_2) = x_2 - \frac{b(x_1x_{14} + x_6x_7 - x_4x_8)}{1 - bx_{12}}, \tag{4.102}$$

$i \in \overline{1,16} \setminus \{2,3,5,11,13\}$,

$$\wp_{12b}(x_3) = x_3 + \frac{b(x_1x_{15} - x_7x_9 + x_4x_{10})}{1 - bx_{12}}, \tag{4.103}$$

$$\wp_{12b}(x_5) = x_5 - \frac{b(x_1x_{16} + x_8x_9 - x_6x_{10})}{1 - bx_{12}}, \tag{4.104}$$

$$\wp_{12b}(x_{11}) = x_{11} + \frac{b(x_4x_{16} + x_6x_{15} + x_9x_{14})}{1 - bx_{12}}, \tag{4.105}$$

$$\wp_{12b}(x_{13}) = x_{13} + \frac{b(x_7x_{16} + x_8x_{15} + x_{10}x_{14})}{1 - bx_{12}}; \tag{4.106}$$

$$\wp_{13b}(x_i) = \frac{x_i}{1 - bx_{13}}, \quad \wp_{13b}(x_1) = x_1 - \frac{b(x_1x_{13} - x_3x_8 + x_5x_7)}{1 - bx_{13}}, \tag{4.107}$$

$i \in \overline{1,16} \setminus \{1,4,6,9,12\}$,

$$\wp_{13b}(x_4) = x_4 + \frac{b(x_2x_{15} + x_3x_{14} + x_7x_{11})}{1 - bx_{13}}, \tag{4.108}$$

$$\wp_{13b}(x_6) = x_6 - \frac{b(x_2x_{16} - x_5x_{14} - x_8x_{11})}{1 - bx_{13}}, \tag{4.109}$$

$$\wp_{13b}(x_9) = x_9 - \frac{b(x_3x_{16} + x_5x_{15} - x_{10}x_{11})}{1 - bx_{13}}, \tag{4.110}$$

$$\wp_{13b}(x_{12}) = x_{12} + \frac{b(x_7x_{16} + x_8x_{15} + x_{10}x_{14})}{1 - bx_{13}}; \tag{4.111}$$

$$\wp_{14b}(x_i) = \frac{x_i}{1 - bx_{14}}, \quad \wp_{14b}(x_1) = x_1 - \frac{b(x_2x_{12} - x_4x_8 + x_6x_7)}{1 - bx_{14}}, \tag{4.112}$$

$i \in \overline{1,16} \setminus \{1,3,5,9,10\}$,

$$\wp_{14b}(x_3) = x_3 - \frac{b(x_2x_{15} - x_4x_{13} + x_7x_{11})}{1 - bx_{14}}, \tag{4.113}$$

$$\wp_{14b}(x_5)=x_5+\frac{b(x_2x_{16}+x_6x_{13}-x_8x_{11})}{1-bx_{14}}, \tag{4.114}$$

$$\wp_{14b}(x_9)=x_9-\frac{b(x_4x_{16}+x_6x_{15}-x_{11}x_{12})}{1-bx_{14}}, \tag{4.115}$$

$$\wp_{14b}(x_{10})=x_{10}-\frac{b(x_7x_{16}+x_8x_{15}-x_{12}x_{13})}{1-bx_{14}}; \tag{4.116}$$

$$\wp_{15b}(x_i)=\frac{x_i}{1-bx_{15}},\quad \wp_{15b}(x_1)=x_1+\frac{b(x_3x_{12}-x_4x_{10}+x_7x_9)}{1-bx_{15}}, \tag{4.117}$$

$i\in\overline{1,16}\setminus\{1,2,5,6,8\}$,

$$\wp_{15b}(x_2)=x_2-\frac{b(x_3x_{14}-x_4x_{13}+x_7x_{11})}{1-bx_{15}}, \tag{4.118}$$

$$\wp_{15b}(x_5)=x_5-\frac{b(x_3x_{16}+x_9x_{13}-x_{10}x_{11})}{1-bx_{15}}, \tag{4.119}$$

$$\wp_{15b}(x_6)=x_6-\frac{b(x_4x_{16}+x_9x_{14}-x_{11}x_{12})}{1-bx_{15}}, \tag{4.120}$$

$$\wp_{15b}(x_8)=x_8-\frac{b(x_7x_{16}+x_{10}x_{14}-x_{12}x_{13})}{1-bx_{15}}; \tag{4.121}$$

$$\wp_{16b}(x_i)=\frac{x_i}{1-bx_{16}},\quad \wp_{16b}(x_1)=x_1-\frac{b(x_5x_{12}-x_6x_{10}+x_8x_9)}{1-bx_{16}}, \tag{4.122}$$

$i\in\overline{1,16}\setminus\{1,2,3,4,7\}$,

$$\wp_{16b}(x_2)=x_2+\frac{b(x_5x_{14}-x_6x_{13}+x_8x_{11})}{1-bx_{16}}, \tag{4.123}$$

$$\wp_{16b}(x_3)=x_3-\frac{b(x_5x_{15}+x_9x_{13}-x_{10}x_{11}))}{1-bx_{16}}, \tag{4.124}$$

$$\wp_{16b}(x_4)=x_4-\frac{b(x_6x_{15}+x_9x_{14}-x_{11}x_{12})}{1-bx_{16}}, \tag{4.125}$$

$$\wp_{16b}(x_7)=x_7-\frac{b(x_8x_{15}+x_{10}x_{14}-x_{12}x_{13})}{1-bx_{16}}; \tag{4.126}$$

where $b\in\mathbb{R}$.

5 Functor from D_5-Mod to E_6-Mod

In this section, we construct a new functor from D_5-**Mod** to E_6-**Mod**.

Note that

$$\begin{aligned} o(10,\mathcal{A}) &= \sum_{1\le p<q\le 5}[\mathcal{A}(E_{p,n+q}-E_{q,n+p})+\mathcal{A}(E_{n+p,q}-E_{n+q,p})] \\ &\quad+\sum_{i,j=1}^{5}\mathcal{A}(E_{i,j}-E_{n+j,n+i}) \end{aligned} \tag{5.1}$$

forms a Lie subalgebra of the matrix algebra $gl(10,\mathcal{A})$ over $\mathcal{A}$ with respect to the commutator, i.e.

$$[fB_1, gB_2] = fg[B_1,B_2] \qquad \text{for } \ f,g\in\mathcal{A},\ B_1,B_2\in gl(10,\mathbb{C}). \tag{5.2}$$

Moreover, we define the Lie algebra

$$\mathcal{K} = o(10,\mathcal{A})\oplus\mathcal{A}\kappa \tag{5.3}$$

with the Lie bracket:

$$[\xi_1+f\kappa,\xi_2+g\kappa]=[\xi_1,\xi_2] \qquad \text{for } \ \xi_1,\xi_2\in o(10,\mathcal{A}),\ f,g\in\mathcal{A}. \tag{5.4}$$

Similarly, $gl(16,\mathcal{A})$ becomes a Lie algebra with the Lie bracket as that in (5.2). Recall the Witt algebra $\mathcal{W}_{16}=\sum_{i=1}^{16}\mathcal{A}\partial_{x_i}$, and Shen [Sg1-3] found a monomorphism $\Im$ from the Lie algebra $\mathcal{W}_{16}$ to the Lie algebra of semi-product $\mathcal{W}_{16}+gl(16,\mathcal{A})$ defined by

$$\Im(\sum_{i=1}^{16} f_i\partial_{x_i})=\sum_{i=1}^{16} f_i\partial_{x_i}+\Im_1(\sum_{i=1}^{16} f_i\partial_{x_i}),\ \ \Im_1(\sum_{i=1}^{16} f_i\partial_{x_i})=\sum_{i,j=1}^{16}\partial_{x_i}(f_j)E_{i,j}. \tag{5.5}$$

According to our construction of P_1-P_{16} in (4.12)-(4.32),

$$\Im_1(P_i)=\sum_{r=1}^{16}x_r\Im_1([\xi_r,\eta_i]|_{\mathcal{A}}) \qquad \text{for } \ i\in\overline{1,16}. \tag{5.6}$$

On the other hand,

$$\widehat{\mathcal{K}}=\mathcal{W}_{16}\oplus\mathcal{K} \tag{5.7}$$

becomes a Lie algebra with the Lie bracket

$$\begin{aligned} & [d_1+f_1B_1+f_2\kappa, d_2+g_1B_2+g_2\kappa] \\ = \ & [d_1,d_2]+f_1g_1[B_1,B_2]+d_1(g_2)B_2-d_2(g_1)B_1+(d_1(g_2)-d_2(g_1))\kappa \end{aligned} \tag{5.8}$$

for $f_1,f_2,g_1,g_2\in\mathcal{A},\ B_1,B_2\in o(10,\mathbb{C})$ and $d_1,d_2\in\mathcal{W}_{16}$. Note

$$\mathcal{G}_0=\mathcal{G}^{D_5}\oplus\mathbb{C}\widehat{\alpha} \tag{5.9}$$

by (2.32) and (4.2). So there exists a Lie algebra monomorphism $\varrho:\mathcal{G}_0\to\mathcal{K}$ determined by

$$\varrho(\widehat{\alpha})=2\kappa,\ \ \varrho(u)=\nu^{-1}(u) \qquad \text{for } \ u\in\mathcal{G}^{D_5}. \tag{5.10}$$

Since $\Im$ is a Lie algebra monomorphism, our construction of P_1-P_{16} in (4.12)-(4.32) and (5.6) shows that we have a Lie algebra monomorphism $\iota:\mathcal{G}^{E_6}\to\widehat{\mathcal{K}}$ given by

$$\iota(u)=u|_{\mathcal{A}}+\varrho(u) \qquad \text{for } \ u\in\mathcal{G}_0, \tag{5.11}$$

$$\iota(\xi_i)=\partial_{x_i},\;\;\iota(\eta_i)=P_i+\sum_{r=1}^{16}x_r\varrho([\xi_r,\eta_i])\qquad\text{for }\;i\in\overline{1,16}.\tag{5.12}$$

In order to calculate $\{\iota(P_1),...,\iota(P_{16})\}$ explicitly, we need the more formulas of ν on the positive root vectors of $o(10,\mathbb{C})$ extended from (2.42)-(2.47). We calculate

$$\nu(E_{3,5}-E_{10,8})=E_{\alpha_4+\alpha_5},\;\;\nu(E_{2,5}-E_{10,7})=-E_{\sum_{i=3}^5\alpha_i},\;\;\nu(E_{1,9}-E_{4,6})=E_{\sum_{i=1}^5\alpha_i},\tag{5.13}$$

$$\nu(E_{3,9}-E_{4,8})=E_{\alpha_2+\alpha_4+\alpha_5},\;\;\nu(E_{2,9}-E_{4,7})=-E_{\sum_{i=2}^5\alpha_i},\tag{5.14}$$

$$\nu(E_{1,5}-E_{10,6})=E_{\alpha_1+\sum_{i=3}^5\alpha_i},\;\;\nu(E_{2,8}-E_{3,7})=-E_{\alpha_4+\sum_{i=2}^5\alpha_i},\tag{5.15}$$

$$\nu(E_{1,8}-E_{3,6})=E_{\alpha_4+\sum_{i=1}^5\alpha_i},\;\;\nu(E_{1,7}-E_{2,6})=-E_{\alpha_3+\alpha_4+\sum_{i=1}^5\alpha_i}.\tag{5.16}$$

Now

$$\begin{aligned}\iota(\eta_1)&=P_1-x_1\varrho(\alpha_6)-x_2\varrho(E_{\alpha_5})-x_3\varrho(E_{\alpha_4+\alpha_5})-x_4\varrho(E_{\alpha_3+\alpha_4+\alpha_5})\\&\quad-x_5\varrho(E_{\alpha_2+\alpha_4+\alpha_5})-x_6\varrho(E_{\sum_{i=2}^5\alpha_i})-x_7\varrho(E_{\alpha_1+\sum_{i=3}^5\alpha_i})-x_8\varrho(E_{\sum_{i=1}^5\alpha_i})\\&\quad-x_9\varrho(E_{\alpha_4+\sum_{i=2}^5\alpha_i})-x_{10}\varrho(E_{\alpha_4+\sum_{i=1}^5\alpha_i})-x_{12}\varrho(E_{\alpha_3+\alpha_4+\sum_{i=1}^5\alpha_i})\\&=P_1+\frac{x_1}{2}[\sum_{i=1}^4(E_{i,i}-E_{5+i,5+i})-E_{5,5}+E_{10,10}-\kappa]-x_2(E_{4,5}-E_{10,9})\\&\quad-x_3(E_{3,5}-E_{10,8})+x_4(E_{2,5}-E_{10,7})-x_5(E_{3,9}-E_{4,8})+x_6(E_{2,9}-E_{4,7})\\&\quad-x_7(E_{1,5}-E_{10,6})-x_8(E_{1,9}-E_{4,6})+x_9(E_{2,8}-E_{3,7})\\&\quad-x_{10}(E_{1,8}-E_{3,6})+x_{12}(E_{1,7}-E_{2,6})\end{aligned}\tag{5.17}$$

by (2.43), (2.46), (2.47), (4.2), (5.10) and (5.13)-(5.16). Moreover,

$$\begin{aligned}\iota(\eta_2)&=\iota([E_{-\alpha_5},\eta_1])=[\iota(E_{-\alpha_5}),\iota(\eta_1)]=-[(E_{5,4}-E_{9,10})|_{\mathcal{A}}+(E_{5,4}-E_{9,10}),\iota(\eta_1)]\\&=-[-x_2\partial_{x_1}+x_{11}\partial_{x_9}+x_{13}\partial_{x_{10}}+x_{14}\partial_{x_{12}}+(E_{5,4}-E_{9,10}),\iota(\eta_1)]\\&=P_2+\frac{x_2}{2}[\sum_{i\neq4}(E_{i,i}-E_{5+i,5+i})-E_{4,4}+E_{9,9}-\kappa]-x_1(E_{5,4}-E_{9,10})\\&\quad-x_3(E_{3,4}-E_{9,8})+x_4(E_{2,4}-E_{9,7})+x_5(E_{3,10}-E_{5,8})-x_6(E_{2,10}-E_{5,7})\\&\quad-x_7(E_{1,4}-E_{9,6})+x_8(E_{1,10}-E_{5,6})-x_{11}(E_{2,8}-E_{3,7})\\&\quad+x_{13}(E_{1,8}-E_{3,6})-x_{14}(E_{1,7}-E_{2,6})\end{aligned}\tag{5.18}$$

by (3.12). According to (3.11),

$$\begin{aligned}\iota(\eta_3)&=\iota([E_{-\alpha_4},\eta_2])=[\iota(E_{-\alpha_4}),\iota(\eta_2)]=-[(E_{4,3}-E_{8,9})|_{\mathcal{A}}+(E_{4,3}-E_{8,9}),\iota(\eta_2)]\\&=-[-x_3\partial_{x_2}-x_9\partial_{x_6}-x_{10}\partial_{x_8}+x_{15}\partial_{x_{14}}+(E_{4,3}-E_{8,9}),\iota(\eta_2)]\\&=P_3+\frac{x_3}{2}[\sum_{i\neq3}(E_{i,i}-E_{5+i,5+i})-E_{3,3}+E_{8,8}-\kappa]-x_1(E_{5,3}-E_{8,10})\\&\quad-x_2(E_{4,3}-E_{8,9})+x_4(E_{2,3}-E_{8,7})-x_5(E_{4,10}-E_{5,9})-x_7(E_{1,3}-E_{8,6})\\&\quad-x_9(E_{2,10}-E_{5,7})+x_{10}(E_{1,10}-E_{5,6})+x_{11}(E_{2,9}-E_{4,7})\\&\quad-x_{13}(E_{1,9}-E_{4,6})+x_{15}(E_{1,7}-E_{2,6}).\end{aligned}\tag{5.19}$$

Furthermore, (3.10) implies

$$\begin{aligned}\iota(\eta_4) &= -\iota([E_{-\alpha_3},\eta_3]) = -[\iota(E_{-\alpha_3}),\iota(\eta_3)] = [(E_{3,2}-E_{7,8})|_{\mathcal{A}} + (E_{3,2}-E_{7,8}),\iota(\eta_3)] \\ &= [x_4\partial_{x_3} + x_6\partial_{x_5} + x_{12}\partial_{x_{10}} + x_{14}\partial_{x_{13}} + (E_{3,2}-E_{7,8}),\iota(\eta_3)] \\ &= P_4 + \frac{x_4}{2}[\sum_{i\neq 2}(E_{i,i}-E_{5+i,5+i}) - E_{2,2} + E_{7,7} - \kappa] + x_1(E_{5,2}-E_{7,10}) \\ &\quad +x_2(E_{4,2}-E_{7,9}) + x_3(E_{3,2}-E_{7,8}) - x_6(E_{4,10}-E_{5,9}) + x_7(E_{1,2}-E_{7,6}) \\ &\quad -x_9(E_{3,10}-E_{5,8}) + x_{11}(E_{3,9}-E_{4,8}) + x_{12}(E_{1,10}-E_{5,6}) \\ &\quad -x_{14}(E_{1,9}-E_{4,6}) + x_{15}(E_{1,8}-E_{3,6}). \end{aligned} \tag{5.20}$$

Observe

$$\begin{aligned}\iota(\eta_5) &= \iota([E_{-\alpha_2},\eta_3]) = [\iota(E_{-\alpha_2}),\iota(\eta_3)] = -[(E_{10,4}-E_{9,5})|_{\mathcal{A}} + (E_{10,4}-E_{9,5}),\iota(\eta_3)] \\ &= -[-x_5\partial_{x_3} - x_6\partial_{x_4} - x_8\partial_{x_7} + x_{16}\partial_{x_{15}} + E_{10,4} - E_{9,5},\iota(\eta_3)] \\ &= P_5 + \frac{x_5}{2}[\sum_{i=1}^{2}(E_{i,i}-E_{5+i,5+i}) - \sum_{i=3}^{5}(E_{i,i}-E_{5+i,5+i}) - \kappa] - x_1(E_{9,3}-E_{8,4}) \\ &\quad +x_2(E_{10,3}-E_{8,5}) - x_3(E_{10,4}-E_{9,5}) + x_6(E_{2,3}-E_{8,7}) - x_8(E_{1,3}-E_{8,6}) \\ &\quad -x_9(E_{2,4}-E_{9,7}) + x_{10}(E_{1,4}-E_{9,6}) - x_{11}(E_{2,5}-E_{10,7}) \\ &\quad +x_{13}(E_{1,5}-E_{10,6}) - x_{16}(E_{1,7}-E_{2,6}) \end{aligned} \tag{5.21}$$

by (3.13). Similarly,

$$\begin{aligned}\iota(\eta_6) &= \iota([E_{-\alpha_2},\eta_4]) = [\iota(E_{-\alpha_2}),\iota(\eta_3)] = -[(E_{10,4}-E_{9,5})|_{\mathcal{A}} + (E_{10,4}-E_{9,5}),\iota(\eta_4)] \\ &= -[-x_5\partial_{x_3} - x_6\partial_{x_4} - x_8\partial_{x_7} + x_{16}\partial_{x_{15}} + E_{10,4} - E_{9,5},\iota(\eta_4)] \\ &= P_6 + \frac{x_6}{2}[\sum_{i=1,3}(E_{i,i}-E_{5+i,5+i}) - \sum_{i=2,4,5}(E_{i,i}-E_{5+i,5+i}) - \kappa] + x_1(E_{9,2}-E_{7,4}) \\ &\quad -x_2(E_{10,2}-E_{7,5}) - x_4(E_{10,4}-E_{9,5}) + x_5(E_{3,2}-E_{7,8}) + x_8(E_{1,2}-E_{7,6}) \\ &\quad -x_9(E_{3,4}-E_{9,8}) - x_{11}(E_{3,5}-E_{10,8}) + x_{12}(E_{1,4}-E_{9,6}) \\ &\quad +x_{14}(E_{1,5}-E_{10,6}) - x_{16}(E_{1,8}-E_{3,6}). \end{aligned} \tag{5.22}$$

Moreover, (3.9) yields

$$\begin{aligned}\iota(\eta_7) &= -\iota([E_{-\alpha_1},\eta_4]) = -[\iota(E_{-\alpha_1}),\iota(\eta_4)] = [(E_{2,1}-E_{6,7})|_{\mathcal{A}} + (E_{2,1}-E_{6,7}),\iota(\eta_4)] \\ &= [x_7\partial_{x_4} + x_8\partial_{x_6} + x_{10}\partial_{x_9} + x_{13}\partial_{x_{11}} + (E_{2,1}-E_{6,7}),\iota(\eta_4)] \\ &= P_7 + \frac{x_7}{2}[\sum_{i2}^{5}(E_{i,i}-E_{5+i,5+i}) - E_{1,1} + E_{6,6} - \kappa] - x_1(E_{5,1}-E_{6,10}) \\ &\quad -x_2(E_{4,1}-E_{6,9}) - x_3(E_{3,1}-E_{6,8}) + x_4(E_{2,1}-E_{6,7}) - x_8(E_{4,10}-E_{5,9}) \\ &\quad -x_{10}(E_{3,10}-E_{5,8}) + x_{12}(E_{2,10}-E_{5,7}) + x_{13}(E_{3,9}-E_{4,8}) \\ &\quad -x_{14}(E_{2,9}-E_{4,7}) + x_{15}(E_{2,8}-E_{3,7}). \end{aligned} \tag{5.23}$$

Furthermore, (3.13) gives

$$\begin{aligned}\iota(\eta_8) &= \iota([E_{-\alpha_2},\eta_7]) = [\iota(E_{-\alpha_2}),\iota(\eta_7)] = -[(E_{10,4}-E_{9,5})|_{\mathcal{A}}+(E_{10,4}-E_{9,5}),\iota(\eta_7)]\\ &= -[-x_5\partial_{x_3}-x_6\partial_{x_4}-x_8\partial_{x_7}+x_{16}\partial_{x_{15}}+E_{10,4}-E_{9,5},\iota(\eta_7)]\\ &= P_8+\frac{x_8}{2}[\sum_{i=2,3}(E_{i,i}-E_{5+i,5+i})-\sum_{i=1,4,5}(E_{i,i}-E_{5+i,5+i})-\kappa]-x_1(E_{9,1}-E_{6,4})\\ &\quad +x_2(E_{10,1}-E_{6,5})-x_5(E_{3,1}-E_{6,8})+x_6(E_{2,1}-E_{6,7})-x_7(E_{10,4}-E_{9,5})\\ &\quad -x_{10}(E_{3,4}-E_{9,8})+x_{12}(E_{2,4}-E_{9,7})-x_{13}(E_{3,5}-E_{10,8})\\ &\quad +x_{14}(E_{2,5}-E_{10,7})-x_{16}(E_{2,8}-E_{3,7}).\end{aligned}\tag{5.24}$$

According to (3.11),

$$\begin{aligned}\iota(\eta_9) &= \iota([E_{-\alpha_4},\eta_6]) = [\iota(E_{-\alpha_4}),\iota(\eta_6)] = -[(E_{4,3}-E_{8,9})|_{\mathcal{A}}+(E_{4,3}-E_{8,9}),\iota(\eta_6)]\\ &= -[-x_3\partial_{x_2}-x_9\partial_{x_6}-x_{10}\partial_{x_8}+x_{15}\partial_{x_{14}}+(E_{4,3}-E_{8,9}),\iota(\eta_6)]\\ &= P_9+\frac{x_9}{2}[\sum_{i=1,4}(E_{i,i}-E_{5+i,5+i})-\sum_{i=2,3,5}(E_{i,i}-E_{5+i,5+i})-\kappa]+x_1(E_{8,2}-E_{7,3})\\ &\quad -x_3(E_{10,2}-E_{7,5})-x_4(E_{10,3}-E_{8,5})-x_5(E_{4,2}-E_{7,9})-x_6(E_{4,3}-E_{8,9}))\\ &\quad +x_{10}(E_{1,2}-E_{7,6})+x_{11}(E_{4,5}-E_{10,9})+x_{12}(E_{1,3}-E_{8,6})\\ &\quad -x_{15}(E_{1,5}-E_{10,6})+x_{16}(E_{1,9}-E_{4,6})\end{aligned}\tag{5.25}$$

and

$$\begin{aligned}\iota(\eta_{10}) &= \iota([E_{-\alpha_4},\eta_8]) = [\iota(E_{-\alpha_4}),\iota(\eta_8)] = -[(E_{4,3}-E_{8,9})|_{\mathcal{A}}+(E_{4,3}-E_{8,9}),\iota(\eta_6)]\\ &= -[-x_3\partial_{x_2}-x_9\partial_{x_6}-x_{10}\partial_{x_8}+x_{15}\partial_{x_{14}}+(E_{4,3}-E_{8,9}),\iota(\eta_8)]\\ &= P_{10}+\frac{x_{10}}{2}[\sum_{i=2,4}(E_{i,i}-E_{5+i,5+i})-\sum_{i=1,3,5}(E_{i,i}-E_{5+i,5+i})-\kappa]-x_1(E_{8,1}-E_{6,3})\\ &\quad +x_3(E_{10,1}-E_{6,5})+x_5(E_{4,1}-E_{6,9})-x_7(E_{10,3}-E_{8,5})-x_8(E_{4,3}-E_{8,9})\\ &\quad +x_9(E_{2,1}-E_{6,7})+x_{12}(E_{2,3}-E_{8,7})+x_{13}(E_{4,5}-E_{10,9})\\ &\quad -x_{15}(E_{2,5}-E_{10,7})+x_{16}(E_{2,9}-E_{4,7}).\end{aligned}\tag{5.26}$$

Moreover, (3.12) yields

$$\begin{aligned}\iota(\eta_{11}) &= -\iota([E_{-\alpha_5},\eta_9]) = -[\iota(E_{-\alpha_5}),\iota(\eta_9)] = -[(E_{5,4}-E_{9,10})|_{\mathcal{A}}+(E_{5,4}-E_{9,10}),\iota(\eta_1)]\\ &= [-x_2\partial_{x_1}+x_{11}\partial_{x_9}+x_{13}\partial_{x_{10}}+x_{14}\partial_{x_{12}}+(E_{5,4}-E_{9,10}),\iota(\eta_9)]\\ &= P_{11}+\frac{x_{11}}{2}[\sum_{i=1,5}(E_{i,i}-E_{5+i,5+i})-\sum_{i=2,3,4}(E_{i,i}-E_{5+i,5+i})-\kappa]-x_2(E_{8,2}-E_{7,3})\\ &\quad +x_3(E_{9,2}-E_{7,4})+x_4(E_{9,3}-E_{8,4})-x_5(E_{5,2}-E_{7,10})-x_6(E_{5,3}-E_{8,10}))\\ &\quad +x_9(E_{5,4}-E_{9,10})+x_{13}(E_{1,2}-E_{7,6})+x_{14}(E_{1,3}-E_{8,6})\\ &\quad +x_{15}(E_{1,4}-E_{9,6})+x_{16}(E_{1,10}-E_{5,6})\end{aligned}\tag{5.27}$$

Furthermore, (3.10) implies

$$\begin{aligned}
\iota(\eta_{12}) &= -\iota([E_{-\alpha_3},\eta_{10}]) = -[\iota(E_{-\alpha_3}),\iota(\eta_{10})] = [(E_{3,2}-E_{7,8})|_{\mathcal{A}} + (E_{3,2}-E_{7,8}),\iota(\eta_3)] \\
&= [x_4\partial_{x_3} + x_6\partial_{x_5} + x_{12}\partial_{x_{10}} + x_{14}\partial_{x_{13}} + (E_{3,2}-E_{7,8}),\iota(\eta_{10})] \\
&= P_{12} + \frac{x_{12}}{2}[\sum_{i=3,4}(E_{i,i}-E_{5+i,5+i}) - \sum_{i=1,2,5}(E_{i,i}-E_{5+i,5+i}) - \kappa] + x_1(E_{7,1}-E_{6,2}) \\
&\quad + x_4(E_{10,1}-E_{6,5}) + x_6(E_{4,1}-E_{6,9}) + x_7(E_{10,2}-E_{7,5}) + x_8(E_{4,2}-E_{7,9}) \\
&\quad + x_9(E_{3,1}-E_{6,8}) + x_{10}(E_{3,2}-E_{7,8}) + x_{14}(E_{4,5}-E_{10,9}) \\
&\quad - x_{15}(E_{3,5}-E_{10,8}) + x_{16}(E_{3,9}-E_{4,8}).
\end{aligned} \tag{5.28}$$

Note that (3.9) gives

$$\begin{aligned}
\iota(\eta_{13}) &= -\iota([E_{-\alpha_1},\eta_{11}]) = -[\iota(E_{-\alpha_1}),\iota(\eta_{11})] = [(E_{2,1}-E_{6,7})|_{\mathcal{A}} + (E_{2,1}-E_{6,7}),\iota(\eta_4)] \\
&= [x_7\partial_{x_4} + x_8\partial_{x_6} + x_{10}\partial_{x_9} + x_{13}\partial_{x_{11}} + (E_{2,1}-E_{6,7}),\iota(\eta_{11})] \\
&= P_{13} + \frac{x_{13}}{2}[\sum_{i=2,5}(E_{i,i}-E_{5+i,5+i}) - \sum_{i=1,3,4}(E_{i,i}-E_{5+i,5+i}) - \kappa] + x_2(E_{8,1}-E_{6,3}) \\
&\quad - x_3(E_{9,1}-E_{6,4}) + x_5(E_{5,1}-E_{6,10}) + x_7(E_{9,3}-E_{8,4}) - x_8(E_{5,3}-E_{8,10})) \\
&\quad + x_{10}(E_{5,4}-E_{9,10}) + x_{11}(E_{2,1}-E_{6,7}) + x_{14}(E_{2,3}-E_{8,7}) \\
&\quad + x_{15}(E_{2,4}-E_{9,7}) + x_{16}(E_{2,10}-E_{5,7}).
\end{aligned} \tag{5.29}$$

Moreover, (3.10) yields

$$\begin{aligned}
\iota(\eta_{14}) &= -\iota([E_{-\alpha_3},\eta_{13}]) = -[\iota(E_{-\alpha_3}),\iota(\eta_{13})] = [(E_{3,2}-E_{7,8})|_{\mathcal{A}} + (E_{3,2}-E_{7,8}),\iota(\eta_{13})] \\
&= [x_4\partial_{x_3} + x_6\partial_{x_5} + x_{12}\partial_{x_{10}} + x_{14}\partial_{x_{13}} + (E_{3,2}-E_{7,8}),\iota(\eta_{13})] \\
&= P_{14} + \frac{x_{14}}{2}[\sum_{i=3,5}(E_{i,i}-E_{5+i,5+i}) - \sum_{i=1,2,4}(E_{i,i}-E_{5+i,5+i}) - \kappa] - x_2(E_{7,1}-E_{6,2}) \\
&\quad - x_4(E_{9,1}-E_{6,4}) + x_6(E_{5,1}-E_{6,10}) - x_7(E_{9,2}-E_{7,4}) + x_8(E_{5,2}-E_{7,10})) \\
&\quad + x_{11}(E_{3,1}-E_{6,8}) + x_{12}(E_{5,4}-E_{9,10}) + x_{13}(E_{3,2}-E_{7,8}) \\
&\quad + x_{15}(E_{3,4}-E_{9,8}) + x_{16}(E_{3,10}-E_{5,8}).
\end{aligned} \tag{5.30}$$

Furthermore, (3.11) implies

$$\begin{aligned}
\iota(\eta_{15}) &= -\iota([E_{-\alpha_4},\eta_{14}]) = -[\iota(E_{-\alpha_4}),\iota(\eta_{14})] = [(E_{4,3}-E_{8,9})|_{\mathcal{A}} + (E_{4,3}-E_{8,9}),\iota(\eta_{14})] \\
&= [-x_3\partial_{x_2} - x_9\partial_{x_6} - x_{10}\partial_{x_8} + x_{15}\partial_{x_{14}} + (E_{4,3}-E_{8,9}),\iota(\eta_{14})] \\
&= P_{15} + \frac{x_{15}}{2}[\sum_{i=4,5}(E_{i,i}-E_{5+i,5+i}) - \sum_{i=1,2,3}(E_{i,i}-E_{5+i,5+i}) - \kappa] + x_3(E_{7,1}-E_{6,2}) \\
&\quad + x_4(E_{8,1}-E_{6,3}) + x_7(E_{8,2}-E_{7,3}) - x_9(E_{5,1}-E_{6,10}) - x_{10}(E_{5,2}-E_{7,10})) \\
&\quad + x_{11}(E_{4,1}-E_{6,9}) - x_{12}(E_{5,3}-E_{8,10}) + x_{13}(E_{4,2}-E_{7,9}) \\
&\quad + x_{14}(E_{4,3}-E_{8,9}) + x_{16}(E_{4,10}-E_{5,9}).
\end{aligned} \tag{5.31}$$

Finally, (3.13) gives

$$\begin{aligned}\iota(\eta_{16}) &= -\iota([E_{-\alpha_2},\eta_{15}]) = -[\iota(E_{-\alpha_2}),\iota(\eta_{15})] = [(E_{10,4}-E_{9,5})|_{\mathcal{A}}+(E_{10,4}-E_{9,5}),\iota(\eta_{15})] \\ &= [-x_5\partial_{x_3}-x_6\partial_{x_4}-x_8\partial_{x_7}+x_{16}\partial_{x_{15}}+E_{10,4}-E_{9,5},\iota(\eta_{15})] \\ &= P_{16}-\frac{x_{16}}{2}[\sum_{i=1}^{5}(E_{i,i}-E_{5+i,5+i})+\kappa]-x_5(E_{7,1}-E_{6,2})-x_6(E_{8,1}-E_{6,3}) \\ &\quad -x_8(E_{8,2}-E_{7,3})+x_9(E_{9,1}-E_{6,4})+x_{10}(E_{9,2}-E_{7,4}))+x_{11}(E_{10,1}-E_{6,5}) \\ &\quad +x_{12}(E_{9,3}-E_{8,4})+x_{13}(E_{10,2}-E_{7,5})+x_{14}(E_{10,3}-E_{8,5}) \\ &\quad +x_{15}(E_{10,4}-E_{9,5}). \end{aligned} \tag{5.32}$$

Recall $\mathcal{A}=\mathbb{C}[x_1,...,x_{16}]$. Let M be an $o(10,\mathbb{C})$-module and set

$$\widehat{M}=\mathcal{A}\otimes_{\mathbb{C}}M. \tag{5.33}$$

We identify

$$f\otimes v=fv \qquad \text{for } f\in\mathcal{A},\ v\in M. \tag{5.34}$$

Recall the Lie algebra $\widehat{\mathcal{K}}$ defined via (5.1)-(5.8). Fix $c\in\mathbb{C}$. Then $\widehat{M}$ becomes a $\widehat{\mathcal{K}}$-module with the action defined by

$$d(fv)=d(f)v,\ \ \kappa|(fv)=cfv,\ \ (gB)(fv)=fgB(v) \tag{5.35}$$

for $f,g\in\mathcal{A},\ v\in M$ and $B\in o(10,\mathbb{C})$.

Since the linear map $\iota:\mathcal{G}^{E_6}\to\widehat{\mathcal{K}}$ defined in (5.10)-(5.12) is a Lie algebra monomorphism, $\widehat{M}$ becomes a $\mathcal{G}^{E_6}$-module with the action defined by

$$\xi(w)=\iota(\xi)(w) \qquad \text{for } \xi\in\mathcal{G}^{E_6},\ w\in\widehat{M}. \tag{5.36}$$

In fact, we have:

Theorem 5.1. *The map $M\mapsto\widehat{M}$ gives a functor from $o(10,\mathbb{C})$-**Mod** to $\mathcal{G}^{E_6}$-**Mod**.*

We remark that the module $\widehat{M}$ is not a generalized module in general because it may not be equal to $U(\mathcal{G})(M)=U(\mathcal{G}_-)(M)$.

Proposition 5.2. *If M is an irreducible $o(10,\mathbb{C})$-module, then $U(\mathcal{G}_-)(M)$ is an irreducible $\mathcal{G}^{E_6}$-module.*

Proof. Note that for any $i\in\overline{1,16}$, $f\in\mathcal{A}$ and $v\in M$, (5.12), (5.35) and (5.36) imply

$$\xi_i(fv)=\partial_{x_i}(f)v. \tag{5.37}$$

Let W be any nonzero $\mathcal{G}^{E_6}$-submodule. The above expression shows that $W\bigcap M\neq\{0\}$. According to (5.35), $W\bigcap M$ is an $o(10,\mathbb{C})$-submodule. By the irreducibility of M, $M\subset W$. Thus $U(\mathcal{G}_-)(M)\subset W$. So $U(\mathcal{G}_-)(M)=W$ is irreducible. $\square$

By the above proposition, the map $M \mapsto U(\mathcal{G}_-)(M)$ is a polynomial extension from irreducible $o(10,\mathbb{C})$-modules to irreducible $\mathcal{G}^{E_6}$-modules.

6 Irreducibility

In this section, we want to determine the irreducibility of $\mathcal{G}^{E_6}$-modules.

Let M be an $o(10,\mathbb{C})$-module and let $\widehat{M}$ be the $\mathcal{G}^{E_6}$-module defined in (5.33)-(5.36). Moreover, $\widehat{M}$ can be viewed as an $o(10,\mathbb{C})$-module with the representation $\iota(\nu(B))|_{\widehat{M}}$ (cf. (2.42)-(2.47)). Indeed, (5.11) and (5.36) show

$$\nu(B)(fv) = B(f)v + fB(v) \qquad \text{for } \ B \in o(10,\mathbb{C}),\ f \in \mathcal{A},\ v \in M \tag{6.1}$$

(cf. (2.49)-(2.71)). So $\widehat{M} = \mathcal{A} \otimes_{\mathbb{C}} M$ is a tensor module of $o(10,\mathbb{C})$. Write

$$\eta^\alpha = \prod_{i=1}^{16} \eta_i^{\alpha_i},\ \ |\alpha| = \sum_{i=1}^{16} \alpha_i \ \text{ for } \ \alpha = (\alpha_1, \alpha_2, ..., \alpha_{16}) \in \mathbb{N}^{16} \tag{6.2}$$

(cf. (2.27)-(2.31)). Recall the Lie subalgebras $\mathcal{G}_\pm$ and $\mathcal{G}_0$ of $\mathcal{G}^{E_6}$ defined in (2.32). For $k \in \mathbb{N}$, we set

$$\mathcal{A}_k = \text{Span}_{\mathbb{C}}\{x^\alpha \mid \alpha \in \mathbb{N}^{16}; |\alpha| = k\},\ \ \widehat{M}_{\langle k\rangle} = \mathcal{A}_k M \tag{6.3}$$

(cf. (2.39), (5.34)) and

$$(U(\mathcal{G}_-)(M))_{\langle k\rangle} = \text{Span}_{\mathbb{C}}\{\eta^\alpha(M) \mid \alpha \in \mathbb{N}^{16},\ |\alpha| = k\}. \tag{6.4}$$

Moreover,

$$(U(\mathcal{G}_-)(M))_{\langle 0\rangle} = \widehat{M}_{\langle 0\rangle} = M. \tag{6.5}$$

Furthermore,

$$\widehat{M} = \bigoplus_{k=0}^{\infty} \widehat{M}_{\langle k\rangle}, \qquad U(\mathcal{G}_-)(M) = \bigoplus_{k=0}^{\infty} (U(\mathcal{G}_-)(M))_{\langle k\rangle}. \tag{6.6}$$

Next we define a linear transformation φ on $\widehat{M}$ determined by

$$\varphi(x^\alpha v) = \eta^\alpha(v) \qquad \text{for } \ \alpha \in \mathbb{N}^{16},\ v \in M. \tag{6.7}$$

Note that $\mathcal{A}_1 = \sum_{i=1}^{16} \mathbb{C}x_i$ forms the 16-dimensional $\mathcal{G}_0$-module (equivalently, the $o(10,\mathbb{C})$ spin module). According to (2.9) and (2.10), $\mathcal{G}_-$ forms a $\mathcal{G}_0$-module with respect to the adjoint representation, and the linear map from $\mathcal{A}_1$ to $\mathcal{G}_0$ determined by $x_i \mapsto \eta_i$ for $i \in \overline{1,16}$ gives a $\mathcal{G}_0$-module isomorphism. Thus φ can also be viewed as a $\mathcal{G}_0$-module homomorphism from $\widehat{M}$ to $U(\mathcal{G}_0)(M)$. Moreover,

$$\varphi(\widehat{M}_{\langle k\rangle}) = (U(\mathcal{G}_-)(M))_{\langle k\rangle} \qquad \text{for } \ k \in \mathbb{N}. \tag{6.8}$$

Note that the Casimir element of $o(10,\mathbb{C})$ is

$$\begin{aligned}\omega &= \sum_{1\leq i<j\leq 10}[(E_{i,5+j}-E_{j,5+i})(E_{5+j,i}-E_{5+i,j})+(E_{5+j,i}-E_{5+i,j})(E_{i,5+j}-E_{j,5+i})] \\ &\quad +\sum_{i,j=1}^{5}(E_{i,j}-E_{5+j,5+i})(E_{j,i}-E_{5+i,5+j})\in U(o(10,\mathbb{C})). \qquad (6.9)\end{aligned}$$

The algebra $U(o(10,\mathbb{C}))$ can be imbedded into the tensor algebra $U(o(10,\mathbb{C}))\otimes U(o(10,\mathbb{C}))$ by the associative algebra homomorphism $\delta : U(o(10,\mathbb{C}))\to U(o(10,\mathbb{C}))\otimes_{\mathbb{C}} U(o(10,\mathbb{C}))$ determined by

$$\delta(u)=u\otimes 1+1\otimes u \qquad \text{for } u\in o(10,\mathbb{C}). \qquad (6.10)$$

Set

$$\tilde{\omega}=\frac{1}{2}(\delta(\omega)-\omega\otimes 1-1\otimes\omega)\in U(o(10,\mathbb{C}))\otimes_{\mathbb{C}} U(o(2n,\mathbb{C})). \qquad (6.11)$$

By (6.9),

$$\begin{aligned}\tilde{\omega} &= \sum_{1\leq i<j\leq 5}[(E_{i,5+j}-E_{j,5+i})\otimes(E_{5+j,i}-E_{5+i,j})+(E_{5+j,i}-E_{5+i,j})\otimes(E_{i,5+j}-E_{j,5+i})] \\ &\quad +\sum_{i,j=1}^{5}(E_{i,j}-E_{5+j,5+i})\otimes(E_{j,i}-E_{5+i,5+j}). \qquad (6.12)\end{aligned}$$

Furthermore, $\tilde{\omega}$ acts on $\widehat{M}$ as an $o(10,\mathbb{C})$-module homomorphism via

$$(B_1\otimes B_2)(fv)=B_1(f)B_2(v) \qquad \text{for } B_1,B_2\in o(10,\mathbb{C}). \qquad (6.13)$$

Lemma 6.1. *We have* $\varphi|_{\widehat{M}_{\langle 1\rangle}}=(\tilde{\omega}-c/2)|_{\widehat{M}_{\langle 1\rangle}}$.

Proof. For any $v\in M$, (2.49)-(2.71), (5.17), (6.12) and (6.13) give

$$\begin{aligned}\tilde{\omega}(x_1v) &= [-x_2(E_{4,5}-E_{10,9})-x_3(E_{3,5}-E_{10,8})+x_4(E_{2,5}-E_{10,7}) \\ &\quad -x_5(E_{3,9}-E_{4,8})-x_7(E_{1,5}-E_{10,6})+x_6(E_{2,9}-E_{4,7}) \\ &\quad -x_8(E_{1,9}-E_{4,6})+x_9(E_{2,8}-E_{3,7})-x_{10}(E_{1,8}-E_{3,6}) \\ &\quad +x_{12}(E_{1,7}-E_{2,6})+x_1(\sum_{i=1}^{4}(E_{i,i}-E_{5+i,5+i})-E_{5,5}+E_{10,10})]v \\ &= \eta_1(v)+(c/2)x_1v=(\varphi+c/2)(x_1v), \qquad (6.14)\end{aligned}$$

equivalently, $\varphi(x_1v)=(\tilde{\omega}-c/2)(x_1v)$. According to (3.12), $(E_{9,10}-E_{5,4})(x_1)=x_2$. So

$$\begin{aligned}&\varphi(x_2v)+\varphi(x_1(E_{9,10}-E_{5,4})(v))=(E_{9,10}-E_{5,4})(\varphi(x_1v) \\ &= (E_{9,10}-E_{5,4})[(\tilde{\omega}-c/2)(x_1v)]=(\tilde{\omega}-c/2)[(E_{9,10}-E_{5,4})(x_1v)] \\ &= (\tilde{\omega}-c/2)[x_2v+x_1(E_{9,10}-E_{5,4})(v)] \\ &= (\tilde{\omega}-c/2)(x_2v)+(\tilde{\omega}-c/2)(x_1(E_{9,10}-E_{5,4})(v)) \\ &= (\tilde{\omega}-c/2)(x_2v)+\varphi(x_1(E_{9,10}-E_{5,4})(v)), \qquad (6.15)\end{aligned}$$

equivalently, $\varphi(x_2v)=(\tilde{\omega}-c/2)(x_2v)$.

Observe that

$$(E_{8,9}-E_{4,3})(x_2)=x_3,\ \ (E_{3,2}-E_{7,8})(x_3)=x_4,\ \ (E_{9,5}-E_{10,4})(x_3)=x_5, \tag{6.16}$$

$$(E_{6,8}-E_{3,1})(x_3)=x_7,\ \ (E_{7,5}-E_{10,2})(x_2)=x_6,\ \ (E_{10,1}-E_{6,5})(x_2)=x_8, \tag{6.17}$$

$$(E_{8,2}-E_{7,3})(x_1)=x_9,\ \ (E_{6,3}-E_{8,1})(x_1)=x_{10},\ \ (E_{7,3}-E_{8,2})(x_2)=x_{11}, \tag{6.17}$$

$$(E_{3,2}-E_{7,8})(x_{10})=x_{12},\ \ (E_{8,1}-E_{6,3})(x_2)=x_{13},\ \ (E_{3,2}-E_{7,8})(x_{13})=x_{14}, \tag{6.18}$$

$$(E_{4,3}-E_{8,9})(x_{14})=x_{15},\ \ (E_{10,4}-E_{9,5})(x_{15})=x_{16} \tag{6.19}$$

by (3.9)-(3.13). Using the argument similar to (6.14) and induction, we can prove

$$\varphi(x_iv)=(\tilde{\omega}-c/2)(x_iv)\qquad \text{for }\ i\in\overline{1,16}, \tag{6.20}$$

equivalently, the lemma holds. $\qquad\Box$

We take the subspace

$$\mathcal{H}=\sum_{i=1}^{5}\mathbb{C}(E_{i,i}-E_{n+i,n+i}) \tag{6.21}$$

as a Cartan subalgebra of the Lie algebra $o(10,\mathbb{C})$ and define $\{\varepsilon_1,...,\varepsilon_5\}\subset\mathcal{H}^*$ by

$$\varepsilon_i(E_{j,j}-E_{n+j,n+j})=\delta_{i,j}. \tag{6.22}$$

The inner product $(\cdot,\cdot)$ on the $\mathbb{Q}$-subspace

$$L_{\mathbb{Q}}=\sum_{i=1}^{5}\mathbb{Q}\varepsilon_i \tag{6.23}$$

is given by

$$(\varepsilon_i,\varepsilon_j)=\delta_{i,j}\qquad \text{for }\ i,j\in\overline{1,n}. \tag{6.24}$$

Then the root system of $o(10,\mathbb{C})$ is

$$\Phi_{D_5}=\{\pm\varepsilon_i\pm\varepsilon_j\mid 1\le i<j\le 5\}. \tag{6.25}$$

We take the set of positive roots

$$\Phi_{D_5}^{+}=\{\varepsilon_i\pm\varepsilon_j\mid 1\le i<j\le 5\}. \tag{6.26}$$

In particular,

$$\Pi_{D_5}=\{\varepsilon_1-\varepsilon_2,...,\varepsilon_4-\varepsilon_5,\varepsilon_4+\varepsilon_5\}\ \text{ is the set of positive simple roots.} \tag{6.27}$$

Recall the set of dominate integral weights

$$\Lambda^+ = \{\mu \in L_{\mathbb{Q}} \mid (\varepsilon_4+\varepsilon_5,\mu),(\varepsilon_i-\varepsilon_{i+1},\mu)\in\mathbb{N} \text{ for } i\in\overline{1,4}\}. \tag{6.28}$$

According to (6.24),

$$\Lambda^+ = \{\mu=\sum_{i=1}^5 \mu_i\varepsilon_i \mid \mu_i\in\mathbb{Z}/2;\mu_i-\mu_{i+1},\mu_4+\mu_5\in\mathbb{N}\}. \tag{6.29}$$

Note that if $\mu\in\Lambda^+$, then $\mu_4\geq|\mu_5|$. Denote

$$\rho=\frac{1}{2}\sum_{\nu\in\Phi_{D_5}^+}\nu. \tag{6.30}$$

Then

$$(\rho,\nu)=1 \qquad \text{for }\ \nu\in\Pi_{D_5} \tag{6.31}$$

(e.g., cf. [Hu]). By (6.24),

$$\rho=\sum_{i=1}^4(5-i)\varepsilon_i. \tag{6.32}$$

For any $\mu\in\Lambda^+$, we denote by $V(\mu)$ the finite-dimensional irreducible $o(10,\mathbb{C})$-module with the highest weight μ and have

$$\omega|_{V(\mu)}=(\mu+2\rho,\mu)\mathrm{Id}_{V(\mu)} \tag{6.33}$$

by (6.9).

Let $\mathbb{Z}_2=\mathbb{Z}/2\mathbb{Z}=\{0,1\}$. According to (2.71) and Table 1, the weight set of the $o(10,\mathbb{C})$-module $\mathcal{A}_1$ is

$$\Pi(\mathcal{A}_1)=\{(1/2)\sum_{i=1}^5(-1)^{k_i}\varepsilon_i\mid k_i\in\mathbb{Z}_2,\ \sum_{i=1}^5 k_i=1\}. \tag{6.34}$$

Fix $\lambda\in\Lambda^+$, we define

$$\Upsilon(\lambda)=\{\lambda+\mu\mid\mu\in\Pi(\mathcal{A}_1),\ \lambda+\mu\in\Lambda^+\}. \tag{6.35}$$

Lemma 6.2. *We have*:

$$\mathcal{A}_1\otimes V(\lambda)\cong\bigoplus_{\lambda'\in\Upsilon(\lambda)}V(\lambda'). \tag{6.36}$$

Proof. Note that all the weight subspaces of $\mathcal{A}_1$ are one-dimensional. Thus all the irreducible components of $\mathcal{A}_1\otimes V(\lambda)$ are of multiplicity one. Since

$$\rho+\lambda+\mu\in\Lambda^+ \qquad \text{for }\ \mu\in\Pi(\mathcal{A}_1), \tag{6.37}$$

the tensor theory of finite-dimensional irreducible modules over a finite-dimensional simple Lie algebra (e.g., cf. [Hu]) says that $V(\lambda')$ is a component of $\mathcal{A}_1 \otimes V(\lambda)$ if and only if $\lambda' \in \Upsilon(\lambda)$. $\qquad\Box$

Recall

$$\text{the highest weight of } \mathcal{A}_1 = \frac{1}{2}(\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4 - \varepsilon_5) = \lambda_4, \tag{6.38}$$

the forth fundamental weight of $o(10, \mathbb{C})$, by (2.71) and Table 1. Thus the eigenvalues of $\tilde{\omega}|_{\widehat{V(\lambda)}_{\langle 1\rangle}}$ are

$$\{[(\lambda' + 2\rho, \lambda') - (\lambda + 2\rho, \lambda) - (\lambda_4 + 2\rho, \lambda_4)]/2 \mid \lambda' \in \Upsilon(\lambda)\} \tag{6.39}$$

by (6.11) and (6.13). Define

$$\ell_\omega(\lambda) = \min\{[(\lambda' + 2\rho, \lambda') - (\lambda + 2\rho, \lambda) - (\lambda_4 + 2\rho, \lambda_4)]/2 \mid \lambda' \in \Upsilon(\lambda)\}, \tag{6.40}$$

which will be used to determine the irreducibility of $\widehat{V(\lambda)}$. If $\lambda' = \lambda + \lambda_4 - \alpha \in \Upsilon(\lambda)$ with $\alpha \in \Phi^+_{D_5}$, then

$$(\lambda' + 2\rho, \lambda') - (\lambda + 2\rho, \lambda) - (\lambda_4 + 2\rho, \lambda_4) = 2[(\lambda, \lambda_4) + 1 - (\rho + \lambda + \lambda_4, \alpha)]. \tag{6.41}$$

Recall the differential operators $P_1, ..., P_{16}$ given in (4.16)-(4.32). We also view the elements of $\mathcal{A}$ as the multiplication operators on $\mathcal{A}$. Recall ζ_1 in (3.8). It turns out that we need the following lemma in order to determine the irreducibility of $\widehat{V(\lambda)}$.

Lemma 6.3. *As operators on* $\mathcal{A}$:

$$P_{11}x_1 + P_1x_{11} + P_9x_2 + P_2x_9 - P_6x_3 - P_3x_6 + P_5x_4 + P_4x_5 = \zeta_1(D-6). \tag{6.42}$$

Proof. According to (4.16), (4.18)-(4.22), (4.25) and (4.27), we find that

$$\begin{aligned} & P_{11}x_1 + P_1x_{11} + P_9x_2 + P_2x_9 - P_6x_3 - P_3x_6 + P_5x_4 + P_4x_5 \\ = \; & -6\zeta_1 + x_1P_{11} + x_{11}P_1 + x_2P_9 + x_9P_2 - x_3P_6 - x_6P_3 + x_4P_5 + x_5P_4 \end{aligned} \tag{6.43}$$

and

$$\begin{aligned} & x_1P_{11} + x_{11}P_1 + x_2P_9 + x_9P_2 - x_3P_6 - x_6P_3 + x_4P_5 + x_5P_4 \\ = \; & x_1(x_{11}D - \zeta_1\partial_{x_1} + \zeta_5\partial_{x_7} + \zeta_9\partial_{x_8} + \zeta_8\partial_{x_{10}} - \zeta_7\partial_{x_{12}}) \\ & +x_{11}(x_1D - \zeta_1\partial_{x_{11}} - \zeta_2\partial_{x_{13}} - \zeta_3\partial_{x_{14}} - \zeta_4\partial_{x_{15}} - \zeta_{10}\partial_{x_{16}}) \\ & +x_2(x_9D - \zeta_1\partial_{x_2} + \zeta_4\partial_{x_7} - \zeta_{10}\partial_{x_8} - \zeta_8\partial_{x_{13}} + \zeta_7\partial_{x_{14}}) \\ & +x_9(x_2D - \zeta_1\partial_{x_9} - \zeta_2\partial_{x_{10}} - \zeta_3\partial_{x_{12}} + \zeta_5\partial_{x_{15}} - \zeta_9\partial_{x_{16}}) \\ & -x_3(x_6D + \zeta_1\partial_{x_3} - \zeta_3\partial_{x_7} + \zeta_{10}\partial_{x_{10}} - \zeta_9\partial_{x_{13}} + \zeta_7\partial_{x_{15}}) \\ & -x_6(x_3D + \zeta_1\partial_{x_6} + \zeta_2\partial_{x_8} + \zeta_4\partial_{x_{12}} + \zeta_5\partial_{x_{14}} - \zeta_8\partial_{x_{16}}) \end{aligned}$$

$$
\begin{aligned}
&+x_4(x_5D-\zeta_1\partial_{x_4}-\zeta_2\partial_{x_7}-\zeta_{10}\partial_{x_{12}}-\zeta_9\partial_{x_{14}}-\zeta_8\partial_{x_{15}})\\
&+x_5(x_4D-\zeta_1\partial_{x_5}+\zeta_3\partial_{x_8}-\zeta_4\partial_{x_{10}}-\zeta_5\partial_{x_{13}}+\zeta_7\partial_{x_{16}})\\
=\;&2\zeta_1D-\zeta_1\sum_{i=1,2,3,4,5,6,9,11}x_i\partial_{x_i}+(x_1\zeta_5+x_2\zeta_4+x_3\zeta_3-x_4\zeta_2)\partial_{x_7}\\
&+(x_1\zeta_9-x_2\zeta_{10}-x_6\zeta_2+x_5\zeta_3)\partial_{x_8}+(x_1\zeta_8-x_9\zeta_2-x_3\zeta_{10}-x_5\zeta_4)\partial_{x_{10}}\\
&-(x_1\zeta_7+x_9\zeta_3+x_6\zeta_4+x_4\zeta_{10})\partial_{x_{12}}-(x_{11}\zeta_2+x_2\zeta_8-x_3\zeta_9+x_5\zeta_5)\partial_{x_{13}}\\
&-(x_{11}\zeta_3-x_2\zeta_7+x_6\zeta_5-x_4\zeta_9)\partial_{x_{14}}-(x_{11}\zeta_4-x_9\zeta_5+x_3\zeta_7+x_4\zeta_8)\partial_{x_{15}}\\
&-(x_{11}\zeta_{10}+x_9\zeta_9-x_6\zeta_8-x_5\zeta_7)\partial_{x_{16}}=\zeta_1D
\end{aligned}\tag{6.44}
$$

because

$$
\begin{aligned}
&x_1\zeta_5+x_2\zeta_4+x_3\zeta_3-x_4\zeta_2=x_1(-x_2x_{15}-x_3x_{14}+x_4x_{13}-x_7x_{11})\\
&+x_2(x_1x_{15}-x_3x_{12}+x_4x_{10}-x_7x_9)+x_3(x_1x_{14}+x_2x_{12}-x_4x_8+x_6x_7)\\
&-x_4(x_1x_{13}+x_2x_{10}-x_3x_8+x_5x_7)=-\zeta_1x_7
\end{aligned}\tag{6.45}
$$

by (3.8) and (3.14)-(3.17),

$$
\begin{aligned}
&x_1\zeta_9-x_2\zeta_{10}-x_6\zeta_2+x_5\zeta_3=x_1(x_2x_{16}-x_5x_{14}+x_6x_{13}-x_8x_{11})\\
&-x_2(x_1x_{16}+x_5x_{12}-x_6x_{10}+x_8x_9)-x_6(x_1x_{13}+x_2x_{10}-x_3x_8+x_5x_7)\\
&+x_5(x_1x_{14}+x_2x_{12}-x_4x_8+x_6x_7)=-\zeta_1x_8
\end{aligned}\tag{6.46}
$$

by (3.8), (3.14), (3.15), (3.18) and (3.19),

$$
\begin{aligned}
&x_1\zeta_8-x_9\zeta_2-x_3\zeta_{10}+x_5\zeta_4=x_1(x_3x_{16}+x_5x_{15}+x_9x_{13}-x_{10}x_{11})\\
&-x_9(x_1x_{13}+x_2x_{10}-x_3x_8+x_5x_7)-x_3(x_1x_{16}+x_5x_{12}-x_6x_{10}+x_8x_9)\\
&-x_5(x_1x_{15}-x_3x_{12}-x_4x_{10}-x_7x_9)=-\zeta_1x_{10}
\end{aligned}\tag{6.47}
$$

by (3.8), (3.14), (3.16), (3.18) and (3.20),

$$
\begin{aligned}
&x_1\zeta_7+x_9\zeta_3+x_6\zeta_4+x_4\zeta_{10}=x_1(-x_4x_{16}-x_6x_{15}-x_9x_{14}+x_{11}x_{12})\\
&+x_9(x_1x_{14}+x_2x_{12}-x_4x_8+x_6x_7)+x_6(x_1x_{15}-x_3x_{12}+x_4x_{10}-x_7x_9)\\
&+x_4(x_1x_{16}+x_5x_{12}-x_6x_{10}+x_8x_9)=\zeta_1x_{12}
\end{aligned}\tag{6.48}
$$

by (3.8), (3.15), (3.16), (3.18) and (3.21),

$$
\begin{aligned}
&x_{11}\zeta_2+x_2\zeta_8-x_3\zeta_9+x_5\zeta_5=x_{11}(x_1x_{13}+x_2x_{10}-x_3x_8+x_5x_7)\\
&+x_2(x_3x_{16}+x_5x_{15}+x_9x_{13}-x_{10}x_{11})-x_3(x_2x_{16}-x_5x_{14}+x_6x_{13}-x_8x_{11})\\
&+x_5(-x_2x_{15}-x_3x_{14}+x_4x_{13}-x_7x_{11})=\zeta_1x_{13}
\end{aligned}\tag{6.49}
$$

by (3.8), (3.14), (3.17), (3.19) and (3.20),

$$\begin{aligned}&x_{11}\zeta_3 - x_2\zeta_7 + x_6\zeta_5 - x_4\zeta_9 = x_{11}(x_1x_{14} + x_2x_{12} - x_4x_8 + x_6x_7)\\ &-x_2(-x_4x_{16} - x_6x_{15} - x_9x_{14} + x_{11}x_{12}) + x_6(-x_2x_{15} - x_3x_{14} + x_4x_{13} - x_7x_{11})\\ &-x_4(x_2x_{16} - x_5x_{14} + x_6x_{13} - x_8x_{11}) = \zeta_1x_{14}\end{aligned} \tag{6.50}$$

by (3.8), (3.15), (3.17), (3.19) and (3.21),

$$\begin{aligned}&x_{11}\zeta_4 - x_9\zeta_5 + x_3\zeta_7 + x_4\zeta_8 = x_{11}(x_1x_{15} - x_3x_{12} + x_4x_{10} - x_7x_9)\\ &-x_9(-x_2x_{15} - x_3x_{14} + x_4x_{13} - x_7x_{11}) + x_3(-x_4x_{16} - x_6x_{15} - x_9x_{14} + x_{11}x_{12})\\ &+x_4(x_3x_{16} + x_5x_{15} + x_9x_{13} - x_{10}x_{11}) = \zeta_1x_{15}\end{aligned} \tag{6.51}$$

by (3.8), (3.16), (3.17), (3.20) and (3.21),

$$\begin{aligned}&x_{11}\zeta_{10} + x_9\zeta_9 - x_6\zeta_8 - x_5\zeta_7 = x_{11}(x_1x_{16} + x_5x_{12} - x_6x_{10} + x_8x_9)\\ &+x_9(x_2x_{16} - x_5x_{14} + x_6x_{13} - x_8x_{11}) - x_6(x_3x_{16} + x_5x_{15} + x_9x_{13} - x_{10}x_{11})\\ &-x_5(-x_4x_{16} - x_6x_{15} - x_9x_{14} + x_{11}x_{12}) = \zeta_1x_{16}\end{aligned} \tag{6.52}$$

by (3.8) and (3.18)-(3.21). $\Box$

We define the multiplication

$$f(gv) = (fg)v \qquad \text{for } \ f, g \in \mathcal{A}, \ v \in M. \tag{6.53}$$

Then (5.17)-(5.22), (5.25) and (5.27) gives

$$\begin{aligned}&\sum_{r=1}^{16} x_r\varrho([\xi_r, x_1\eta_{11} + x_{11}\eta_1 + x_2\eta_9 + x_9\eta_2 - x_3\eta_6 - x_6\eta_3 + x_4\eta_5 + x_5\eta_4])\\ =\ & x_1[x_{13}(E_{1,2} - E_{7,6}) + x_{14}(E_{1,3} - E_{8,6}) + x_{15}(E_{1,4} - E_{9,6}) + x_{16}(E_{1,10} - E_{5,6})]\\ &-x_{11}[x_7(E_{1,5} - E_{10,6}) + x_8(E_{1,9} - E_{4,6}) + x_{10}(E_{1,8} - E_{3,6}) - x_{12}(E_{1,7} - E_{2,6})]\\ &+x_2[x_{10}(E_{1,2} - E_{7,6}) + x_{12}(E_{1,3} - E_{8,6}) - x_{15}(E_{1,5} - E_{10,6}) + x_{16}(E_{1,9} - E_{4,6})]\\ &-x_9[x_7(E_{1,4} - E_{9,6}) - x_8(E_{1,10} - E_{5,6}) - x_{13}(E_{1,8} - E_{3,6}) + x_{14}(E_{1,7} - E_{2,6})]\\ &-x_3[x_8(E_{1,2} - E_{7,6}) + x_{12}(E_{1,4} - E_{9,6}) + x_{14}(E_{1,5} - E_{10,6}) - x_{16}(E_{1,8} - E_{3,6})]\\ &+x_6[x_7(E_{1,3} - E_{8,6}) - x_{10}(E_{1,10} - E_{5,6}) + x_{13}(E_{1,9} - E_{4,6}) - x_{15}(E_{1,7} - E_{2,6})]\\ &-x_4[x_8(E_{1,3} - E_{8,6}) - x_{10}(E_{1,4} - E_{9,6}) - x_{13}(E_{1,5} - E_{10,6}) + x_{16}(E_{1,7} - E_{2,6})]\\ &+x_5[x_7(E_{1,2} - E_{7,6}) + x_{12}(E_{1,10} - E_{5,6}) - x_{14}(E_{1,9} - E_{4,6}) + x_{15}(E_{1,8} - E_{3,6})]\\ &+\zeta_1(E_{1,1} - E_{6,6} - c)\\ =\ & \sum_{i=1}^{5}\zeta_i(E_{1,i} - E_{5+i,6}) + \sum_{r=2}^{5}\zeta_{5+r}(E_{1,5+r} - E_{r,6}) - c\zeta_1\end{aligned} \tag{6.54}$$

as operators on $\widehat{M}$ (cf. (5.33)), where ζ_i are defined in (3.8) and (3.14)-(3.22). By Lemma 6.3, (5.5), (5.6) and (6.54),

$$\begin{aligned} T_1 &= \iota(\eta_{11})x_1+\iota(\eta_1)x_{11}+\iota(\eta_9)x_2+\iota(\eta_2)x_9-\iota(\eta_6)x_3-\iota(\eta_3)x_6+\iota(\eta_5)x_4+\iota(\eta_4)x_5 \\ &= \zeta_1(D-c-6)+\sum_{i=1}^{5}\zeta_i(E_{1,i}-E_{5+i,6})+\sum_{r=2}^{5}\zeta_{5+r}(E_{1,5+r}-E_{r,6}) \end{aligned}\tag{6.55}$$

as operators on $\widehat{M}$. We define an $o(10,\mathbb{C})$-module structure on the space $\mathrm{End}\,\widehat{M}$ of linear transformations on $\widehat{M}$ by

$$B(T)=[\nu(B),T]=\nu(B)T-T\nu(B)\qquad\text{for }\ B\in o(10,\mathbb{C}),\ T\in\mathrm{End}\,\widehat{M}\tag{6.56}$$

(cf. (6.1)). It can be verified that T_1 is an $o(10,\mathbb{C})$-singular vector with weight ε_1 in $\mathrm{End}\,\widehat{M}$. So it generates the 10-dimensional natural module.

We set

$$\begin{aligned} T_2 &= -[\iota(E_{-\alpha_1}),T_1]=[(E_{2,1}-E_{6,7})|_{\mathcal{A}}+(E_{2,1}-E_{6,7}),T_1] \\ &= \zeta_2(D-c-6)+\sum_{i=1}^{5}\zeta_i(E_{2,i}-E_{5+i,7})+\sum_{r=1,3,4,5}\zeta_{5+r}(E_{2,5+r}-E_{r,7})], \end{aligned}\tag{6.57}$$

$$\begin{aligned} T_3 &= -[\iota(E_{-\alpha_3}),T_2]=[(E_{3,2}-E_{7,8})|_{\mathcal{A}}+(E_{3,2}-E_{7,8}),T_2] \\ &= \zeta_3(D-c-6)+\sum_{i=1}^{5}\zeta_i(E_{3,i}-E_{5+i,8})+\sum_{r=1,2,4,5}\zeta_{5+r}(E_{3,5+r}-E_{r,8})], \end{aligned}\tag{6.58}$$

$$\begin{aligned} T_4 &= -[\iota(E_{-\alpha_4}),T_3]=[(E_{4,3}-E_{8,9})|_{\mathcal{A}}+(E_{4,3}-E_{8,9}),T_3] \\ &= \zeta_4(D-c-6)+\sum_{i=1}^{5}\zeta_i(E_{4,i}-E_{5+i,9})+\sum_{r=1,2,3,5}\zeta_{5+r}(E_{4,5+r}-E_{r,9})], \end{aligned}\tag{6.59}$$

$$\begin{aligned} T_5 &= -[\iota(E_{-\alpha_5}),T_4]=[(E_{5,4}-E_{9,10})|_{\mathcal{A}}+(E_{5,4}-E_{9,10}),T_4] \\ &= \zeta_5(D-c-6)+\sum_{i=1}^{5}\zeta_i(E_{5,i}-E_{5+i,10})+\sum_{r=1}^{4}\zeta_{5+r}(E_{5,5+r}-E_{r,10}), \end{aligned}\tag{6.60}$$

$$\begin{aligned} T_{10} &= -[\iota(E_{-\alpha_2}),T_4]=[(E_{10,4}-E_{9,5})|_{\mathcal{A}}+(E_{10,4}-E_{9,5}),T_4] \\ &= \zeta_{10}(D-c-6)+\sum_{i=1}^{4}\zeta_i(E_{10,i}-E_{5+i,5})+\sum_{r=1}^{5}\zeta_{5+r}(E_{10,5+r}-E_{r,5}), \end{aligned}\tag{6.61}$$

$$\begin{aligned} T_9 &= -[\iota(E_{-\alpha_2}),T_5]=[(E_{10,4}-E_{9,5})|_{\mathcal{A}}+(E_{10,4}-E_{9,5}),T_5] \\ &= \zeta_9(D-c-6)+\sum_{i=1,2,3,5}\zeta_i(E_{9,i}-E_{5+i,4})+\sum_{r=1}^{5}\zeta_{5+r}(E_{9,5+r}-E_{r,4}), \end{aligned}\tag{6.62}$$

$$\begin{aligned} T_8 &= [\iota(E_{-\alpha_4}), T_9] = [(E_{8,9} - E_{4,3})|_{\mathcal{A}} + (E_{8,9} - E_{4,3}), T_9] \\ &= \zeta_8(D - c - 6) + \sum_{i=1,2,4,5} \zeta_i(E_{8,i} - E_{5+i,3}) + \sum_{r=1}^{5} \zeta_{5+r}(E_{8,5+r} - E_{r,3}), \end{aligned} \tag{6.63}$$

$$\begin{aligned} T_7 &= [\iota(E_{-\alpha_3}), T_8] = [(E_{7,8} - E_{3,2})|_{\mathcal{A}} + (E_{7,8} - E_{3,2}), T_8] \\ &= \zeta_7(D - c - 6) + \sum_{i=1,3,4,5} \zeta_i(E_{7,i} - E_{5+i,2}) + \sum_{r=1}^{5} \zeta_{5+r}(E_{7,5+r} - E_{r,2}), \end{aligned} \tag{6.64}$$

$$\begin{aligned} T_6 &= [\iota(E_{-\alpha_1}), T_7] = [(E_{6,7} - E_{2,1})|_{\mathcal{A}} + (E_{6,7} - E_{2,1}), T_7] \\ &= \zeta_6(D - c - 6) + \sum_{i=2}^{5} \zeta_i(E_{6,i} - E_{5+i,1}) + \sum_{r=1}^{5} \zeta_{5+r}(E_{6,5+r} - E_{r,1}). \end{aligned} \tag{6.65}$$

Then $\mathcal{T} = \sum_{i=1}^{10} \mathbb{C}T_i$ forms the 10-dimensional natural module of $o(10,\mathbb{C})$ with the standard basis $\{T_1, ..., T_{10}\}$.

Denote

$$T_i' = T_i - \zeta_i(D - c - 6) \qquad \text{for } \ i \in \overline{1, 10}. \tag{6.66}$$

Easily see that $\mathcal{T}' = \sum_{i=1}^{10} \mathbb{C}T_i'$ forms the 10-dimensional natural module of $o(10,\mathbb{C})$ with the standard basis $\{T_1', ..., T_{10}'\}$. So we have the $o(10,\mathbb{C})$-module isomorphism from $U = \sum_{i=1}^{10} \mathbb{C}\zeta_i$ to $\mathcal{T}'$ determined by $\zeta_i \mapsto T_i'$ for $i \in \overline{1, 10}$. The weight set of U is

$$\Pi(U) = \{\pm\varepsilon_1, ..., \pm\varepsilon_5\}. \tag{6.67}$$

Let $\lambda \in \Lambda^+$. Denote

$$\Upsilon'(\lambda) = \{\lambda + \mu \mid \mu \in \Pi(U), \ \ \lambda + \mu \in \Lambda^+\}. \tag{6.68}$$

Take $M = V(\lambda)$. It is known that

$$UV(\lambda) = U \otimes_{\mathbb{C}} V(\lambda) \cong \bigoplus_{\lambda' \in \Upsilon'(\lambda)} V(\lambda'). \tag{6.69}$$

Given $\lambda' \in \Upsilon'(\lambda)$, we pick a singular vector

$$u = \sum_{i=1}^{10} \zeta_i u_i \tag{6.70}$$

of weight λ' in $UV(\lambda)$, where $u_i \in V(\lambda)$. Moreover, any singular vector of weight λ' in $UV(\lambda)$ is a scalar multiple of u. Note that the vector

$$w = \sum_{i=1}^{10} T_i'(u_i) \tag{6.71}$$

is also a singular vector of weight λ' if it is not zero. Thus

$$w = \flat_{\lambda'} u, \qquad \flat_{\lambda'} \in \mathbb{C}. \tag{6.72}$$

Set

$$\flat(\lambda) = \min\{\flat_{\lambda'} \mid \lambda' \in \Upsilon'(\lambda)\}. \tag{6.73}$$

Theorem 6.4. *The $\mathcal{G}^{E_6}$-module $\widehat{V(\lambda)}$ is irreducible if*

$$c \in \mathbb{C} \setminus \{\flat(\lambda) - 6 + \mathbb{N}, 2\ell_\omega(\lambda) + 2\mathbb{N}\}. \tag{6.74}$$

Proof. Recall that the $\mathcal{G}^{E_6}$-submodule $U(\mathcal{G}_-)(V(\lambda))$ is irreducible by Proposition 5.2. It is enough to prove $\widehat{V(\lambda)} = U(\mathcal{G}_-)(V(\lambda))$. It is obvious that

$$\widehat{V(\lambda)}_{\langle 0\rangle} = V(\lambda) = (U(\mathcal{G}_-)(V(\lambda)))_{\langle 0\rangle} \tag{6.75}$$

(cf. (6.3) and (6.4) with $M = V(\lambda)$). Moreover, Lemma 6.1 with $M = V(\lambda)$, (6.40) and (6.74) imply that $\varphi|_{\widehat{V(\lambda)}_{\langle 1\rangle}}$ is invertible, equivalently,

$$\widehat{V(\lambda)}_{\langle 1\rangle} = (U(\mathcal{G}_-)(V(\lambda)))_{\langle 1\rangle}. \tag{6.76}$$

Suppose that

$$\widehat{V(\lambda)}_{\langle i\rangle} = (U(\mathcal{G}_-)(V(\lambda)))_{\langle i\rangle} \tag{6.77}$$

for $i \in \overline{0,k}$ with $1 \leq k \in \mathbb{N}$.

For any $v \in V(\lambda)$ and $\alpha \in \mathbb{N}^{16}$ such that $|\alpha| = k-1$, we have

$$T_r(x^\alpha v) = x^\alpha[(|\alpha| - c - 6)\zeta_r + T_r'](v) \in (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle}, \qquad r \in \overline{1,10} \tag{6.78}$$

by (6.77) with $i = k-1, k$. But

$$V' = \mathrm{Span}\{[(|\alpha| - c - 6)\zeta_r + T_r'](v) \mid r \in \overline{1,10},\ v \in V(\lambda)\} \tag{6.79}$$

forms an $o(10,\mathbb{C})$-submodule of $UV(\lambda)$ with respect to the action in (6.1). Let u be a $o(10,\mathbb{C})$-singular in (6.70). Then

$$V' \ni \sum_{r=1}^{10}[(|\alpha| - c - 6)\zeta_r + T_r'](u_r) = (|\alpha| - c - 6)u + w = (|\alpha| - c - 6 + \flat_{\lambda'})u \tag{6.80}$$

by (6.71) and (6.72). Moreover, (6.73) and (6.74) yield $u \in V'$. Since $UV(\lambda)$ is an $o(10,\mathbb{C})$-module generated by all the singular vectors, we have $V' = UV(\lambda)$. So

$$x^\alpha UV(\lambda) \subset (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle}. \tag{6.81}$$

The arbitrariness of α implies

$$\zeta_r\widehat{V(\lambda)}_{\langle k-1\rangle}\subset (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle}\qquad\text{for }\; r\in\overline{1,10}. \tag{6.82}$$

Given any $f\in\mathcal{A}_k$ and $v\in V(\lambda)$, we have

$$\zeta_r\partial_{x_i}(f)v\in\zeta_r\widehat{V(\lambda)}_{\langle k-1\rangle}\subset (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle}\qquad\text{for }\; r\in\overline{1,10},\; i\in\overline{1,16}. \tag{6.83}$$

Moreover,

$$\begin{aligned}\eta_s(fv) &= \iota(\eta_s)(fv)=P_s(fv)+f(\tilde\omega-c/2)(x_sv)\\ &\equiv f(k+\tilde\omega-c/2)(x_sv)\;\;(\mathrm{mod}\;\sum_{r=1}^{10}\zeta_r\widehat{V(\lambda)}_{\langle k-1\rangle})\end{aligned} \tag{6.84}$$

for $s\in\overline{1,16}$ by (4.16)-(4.32), (5.17)-(5.32) and Lemma 6.1. According to (6.40), (6.74), (6.82) and (6.84), we get

$$x_sfv\in (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle}\qquad\text{for }\; s\in\overline{1,16}. \tag{6.85}$$

Thus (6.77) holds for $i=k+1$. By induction on k, (6.77) holds for any $i\in\mathbb{N}$, that is, $\widehat{V(\lambda)}=U(\mathcal{G}_-)(V(\lambda))$. $\qquad\Box$

When $\lambda=0$, $V(0)$ is the one-dimensional trivial module and $\ell_\omega(0)=\flat(0)=0$. So we have:

Corollary 6.5. *The $\mathcal{G}^{E_6}$-module $\widehat{V(0)}$ is irreducible if $c\in\mathbb{C}\setminus\{\mathbb{N}-6\}$.*

Next we consider the case $\lambda=k\varepsilon_1=k\lambda_1$ for some positive integer k, where λ_1 is the first fundamental weight. Note

$$\Upsilon(k\varepsilon_1)=\{\lambda_4+k\varepsilon_1,\lambda_4+(k-1)\varepsilon_1+\varepsilon_5\} \tag{6.86}$$

by (6.34) and (6.35). Thus (6.40) and (6.41) give

$$\ell_\omega(k\varepsilon_1)=-4-k/2. \tag{6.87}$$

In order to calculate $\flat(k\varepsilon_1)$, we give a realization of $V(k\varepsilon_1)$. Observe that we have a representation of $o(10,\mathbb{C})$ on $\mathcal{B}=\mathbb{C}[y_1,...,y_{10}]$ determined via

$$E_{i,j}|_{\mathcal{B}}=y_i\partial_{y_j}\qquad\text{for }\; i,j\in\overline{1,10}. \tag{6.88}$$

Denote by $\mathcal{B}_k$ the subspace of homogenous polynomials in $\mathcal{B}$ with degree k. Set

$$\mathcal{H}_k=\{h\in\mathcal{B}_k\mid(\sum_{i=1}^5\partial_{y_i}\partial_{y_{5+i}})(h)=0\}. \tag{6.89}$$

Then $\mathcal{H}_k \cong V(k\varepsilon_1)$ and y_1^k is a highest-weight vector.

According to (6.67) and (6.68),

$$\Upsilon'(k\varepsilon_1) = \{(k+1)\varepsilon_1, (k-1)\varepsilon_1, k\varepsilon_1 + \varepsilon_2\}. \tag{6.90}$$

The vector $\zeta_1 y_1^k$ is a singular vector in $U\mathcal{H}_k$ with weight $(k+1)\varepsilon_1$, where we take $M = \mathcal{H}_k$ in the earlier settings. By (6.55) and (6.66),

$$T_1'(y_1^k) = k\zeta_1 y_1^k \Longrightarrow \flat_{(k+1)\varepsilon_1} = k. \tag{6.91}$$

Moreover, $\zeta_1 y_1^{k-1} y_2 - \zeta_2 y_1^k$ is a singular vector in $U\mathcal{H}_k$ with weight $k\varepsilon_1 + \varepsilon_2$. By (6.55), (6.57) and (6.66), we find

$$\begin{aligned} T_1'(y_1^{k-1}y_2) - T_2'(y_1^k) &= (k-1)\zeta_1 y_1^{k-1} y_2 + \zeta_2 y_1^k - k\zeta_1 y_1^{k-1} y_2 \\ &= \zeta_2 y_1^k - \zeta_1 y_1^{k-1} y_2 = -(\zeta_1 y_1^{k-1} y_2 - \zeta_2 y_1^k). \end{aligned} \tag{6.92}$$

Thus $\flat_{k\varepsilon_1+\varepsilon_2} = -1$. Furthermore,

$$\varpi = (k+3)\sum_{i=1}^{5}[\zeta_i y_1^{k-1} y_{5+i} + \zeta_{5+i} y_1^{k-1} y_i] - (k-1)\zeta_1 y_1^{k-2} \sum_{s=1}^{5} y_s y_{5+s} \tag{6.93}$$

is a singular vector in $U\mathcal{H}_k$ with weight $(k-1)\varepsilon_1$. Expressions (6.55) and (6.57)-(6.66) yield

$$\begin{aligned} & (k+3)\sum_{i=1}^{5}[T_i'(y_1^{k-1}y_{5+i}) + T_{5+i}'(y_1^{k-1}y_i)] - (k-1)T_1'(x^\alpha y_1^{k-2}\sum_{s=1}^{5} y_s y_{5+s}) \\ = \; & (k+3)\sum_{i=1}^{5}[(k-1)\zeta_1 y_1^{k-2} y_i y_{5+i} - \sum_{r=1}^{5}\zeta_r y_1^{k-1} y_{5+r} - \sum_{s\neq i}\zeta_{5+s} y_1^{k-1} y_s \\ & -((k-1)\delta_{i,1}+1)(\sum_{r\neq i}\zeta_r y_1^{k-1} y_{5+r} + \sum_{s=1}^{5}\zeta_{5+s} y_1^{k-1} y_s) \\ & +(k-1)(1-\delta_{i,1})\zeta_1 y_1^{k-2} y_{5+i} y_i] - (k-1)(k-2)\sum_{i=1}^{5}\zeta_1 y_1^{k-2} y_i y_{5+i} \\ = \; & (-8-k)x^\alpha \varpi \Longrightarrow \flat_{(k-1)\varepsilon_1} = -8-k. \end{aligned} \tag{6.94}$$

Therefore, $\flat(k\varepsilon_1) = -8-k$. By Theorem 6.4 and (6.87), we obtain:

Corollary 6.6. *The $\mathcal{G}^{E_6}$-module $\widehat{V(k\varepsilon_1)}$ is irreducible if $c \in \mathbb{C} \setminus \{\mathbb{N} - 14 - k\}$.*

Now we want to consider the cases $\lambda = \varepsilon_1 + \varepsilon_2 = \lambda_2$ (the second fundamental weight) and $\lambda = \varepsilon_1 + \varepsilon_2 + \varepsilon_3 = \lambda_3$ (the third fundamental weight). Let $\mathcal{E}$ be the associative algebra generated by $\{\theta_1, ..., \theta_{10}\}$ with the defining relations:

$$\theta_i\theta_j = -\theta_j\theta_i \qquad \text{for } \; i,j \in \overline{1,10}. \tag{6.95}$$

Moreover, we define linear operators $\{\partial_{\theta_1},...,\partial_{\theta_{10}}\}$ on $\mathcal{E}$ by

$$\partial_{\theta_i}(1)=0,\ \partial_{\theta_i}(\theta_j w)=\delta_{i,j}w-\theta_j\partial_{\theta_i}(w)\qquad\text{for }\ i,j\in\overline{1,10},\ w\in\mathcal{E}.\tag{6.96}$$

The representation of $o(10,\mathbb{C})$ on $\mathcal{E}$ is defined via

$$E_{i,j}|_{\mathcal{E}}=\theta_i\partial_{\theta_j}\qquad\text{for }\ i,j\in\overline{1,10}.\tag{6.97}$$

Denote

$$\mathcal{E}_r=\text{Span}\{\theta_{i_1}\theta_{i_2}\cdots\theta_{i_r}\mid 1\leq i_1<i_2<\cdots<i_r\leq 10\}.\tag{6.98}$$

Then $\mathcal{E}_2$ forms an irreducible $o(10,\mathbb{C})$-submodule isomorphic $V(\lambda_2)$ with a highest weight vector $\theta_1\theta_2$ and $\mathcal{E}_3$ forms an irreducible $o(10,\mathbb{C})$-submodule isomorphic $V(\lambda_3)$ with a highest weight vector $\theta_1\theta_2\theta_3$.

Note that (6.35) gives

$$\Upsilon(\lambda_2)=\{\lambda_4+\lambda_2,\lambda_4,\lambda_4+\varepsilon_1+\varepsilon_5\},\tag{6.99}$$

which yields $\ell_\omega(\lambda_2)=-8$. Moreover, (6.67) and (6.68) imply

$$\Upsilon'(\lambda_2)=\{\varepsilon_1+\lambda_2,\lambda_3,\varepsilon_1\}.\tag{6.100}$$

We have an $o(10,\mathbb{C})$-singular vector $\zeta_1\theta_1\theta_2$ of weight $\varepsilon_1+\lambda_2$ in $U\mathcal{E}_2$, where we take $M=\mathcal{E}_2$ in the earlier settings. By (6.55) and (6.66),

$$T_1'(\theta_1\theta_2)=\zeta_1\theta_1\theta_2\Longrightarrow \flat_{\varepsilon_1+\lambda_2}=1\tag{6.101}$$

Furthermore,,

$$u=\sum_{i=2}^5\zeta_{5+i}\theta_1\theta_i+\sum_{r=1}^5\zeta_r\theta_1\theta_{5+r}\tag{6.102}$$

is an $o(10,\mathbb{C})$-singular vector of weight ε_1 in $U\mathcal{E}_2$ and

$$w=\zeta_1\theta_2\theta_3-\zeta_2\theta_1\theta_3+\zeta_3\theta_1\theta_2\tag{6.103}$$

is an $o(10,\mathbb{C})$-singular vector of weight λ_3 in $U\mathcal{E}_2$. According to (6.55), (6.57), (6.58) and (6.66), we have

$$\begin{aligned}&T_1'(\theta_2\theta_3)-T_2'(\theta_1\theta_3)+T_3'(\theta_1\theta_2)\\ =\ &\zeta_2\theta_1\theta_3-\zeta_3\theta_1\theta_2-\zeta_1\theta_2\theta_3-\zeta_3\theta_1\theta_2-\zeta_1\theta_2\theta_3+\zeta_2\theta_1\theta_3=-2w.\end{aligned}\tag{6.104}$$

So $\flat_{\lambda_3}=-2$. Expressions (6.55) and (6.57)-(6.66) give rise to

$$\begin{aligned}&\sum_{i=2}^5 T_{5+i}'(\theta_1\theta_i)+\sum_{r=1}^5 T_r'(\theta_1\theta_{5+r})\\ =\ &-[\sum_{i=2}^5\zeta_{5+i}\theta_1\theta_i+\sum_{r=2}^5\zeta_r\theta_1\theta_{5+r}]-\sum_{i=2}^5[\sum_{i\neq r\in\overline{1,5}}\zeta_r\theta_1\theta_{5+r}+\sum_{r=2}^5\zeta_{5+r}\theta_1\theta_r]\\ &-\sum_{r=1}^5[\sum_{i=1}^5\zeta_i\theta_1\theta_{5+i}+\sum_{r\neq i\in\overline{2,5}}\zeta_{5+i}\theta_1\theta_i]=-9u.\end{aligned}\tag{6.105}$$

Hence $\flat_{\varepsilon_1}=-9$. Therefore, $\flat(\lambda_2)=-9$. Theorem 6.4 implies:

Corollary 6.7. *The $\mathcal{G}^{E_6}$-module $\widehat{V(\lambda_2)}$ is irreducible if $c\in\mathbb{C}\setminus\{\mathbb{N}-16\}$.*

Observe that (6.35) gives

$$\Upsilon(\lambda_3)=\{\lambda_4+\lambda_3,\lambda_4+\lambda_2+\varepsilon_5,\lambda_4+\varepsilon_1,\lambda_4+\varepsilon_5\},\tag{6.106}$$

which yields $\ell_\omega(\lambda_3)=-21/2$. Moreover,

$$\Upsilon'(\lambda_2)=\{\varepsilon_1+\lambda_3,\sum_{i=1}^4\varepsilon_i,\lambda_2\}.\tag{6.107}$$

Similarly we have $\flat_{\lambda_4+\lambda_3}=1$. Furthermore,,

$$u=\sum_{i=3}^5\zeta_{5+i}\theta_1\theta_2\theta_i+\sum_{r=1}^5\zeta_r\theta_1\theta_2\theta_{5+r}\tag{6.108}$$

is an $o(10,\mathbb{C})$-singular vector of weight λ_2 in $U\mathcal{E}_3$ and

$$w=\zeta_1\theta_2\theta_3\theta_4-\zeta_2\theta_1\theta_3\theta_4+\zeta_3\theta_1\theta_2\theta_4-\zeta_4\theta_1\theta_2\theta_3\tag{6.109}$$

is an $o(10,\mathbb{C})$-singular vector of weight $\sum_{i=1}^4\varepsilon_i$ in $U\mathcal{E}_3$. According to (6.55), (6.57)-(6.59) and (6.66),

$$\begin{aligned}&T_1'(\theta_2\theta_3\theta_4)-T_2'(\theta_1\theta_3\theta_4)+T_3'(\theta_1\theta_2\theta_4)-T_4'(\theta_1\theta_2\theta_3)\\ =\;&\zeta_2\theta_1\theta_3\theta_4-\zeta_3\theta_1\theta_2\theta_4+\zeta_4\theta_1\theta_2\theta_3-\zeta_1\theta_2\theta_3\theta_4-\zeta_3\theta_1\theta_2\theta_4+\zeta_4\theta_1\theta_2\theta_3-\zeta_1\theta_2\theta_3\theta_4\\ &+\zeta_2\theta_1\theta_3\theta_4+\zeta_4\theta_1\theta_2\theta_3-\zeta_1\theta_2\theta_3\theta_4+\zeta_2\theta_1\theta_3\theta_4-\zeta_3\theta_1\theta_2\theta_4=-3w,\end{aligned}\tag{6.110}$$

equivalently, $\flat_{\sum_{i=1}^4\varepsilon_i}=-3$. Expressions (6.55) and (6.57)-(6.66) give rise to

$$\begin{aligned}&\sum_{i=3}^5T_{5+i}'(\theta_1\theta_2\theta_i)+\sum_{r=1}^5T_r(\theta_1\theta_2\theta_{5+r})\\ =\;&-\sum_{i=3}^5(\zeta_{5+i}\theta_1\theta_2\theta_i+\zeta_i\theta_1\theta_2\theta_{5+i})-\sum_{i=3}^5[\sum_{i\neq r\in\overline{1,5}}\zeta_r\theta_1\theta_2\theta_{5+r}+\sum_{r=3}^5\zeta_{5+r}\theta_1\theta_2\theta_r]\\ &-\sum_{r=1}^5[\sum_{i=1}^5\zeta_i\theta_1\theta_2\theta_{5+i}+\sum_{r\neq i\in\overline{3,5}}\zeta_{5+i}\theta_1\theta_2\theta_i]=-8u.\end{aligned}\tag{6.111}$$

So $\flat_{\lambda_2}=-8$. Therefore, $\flat(\lambda_3)=-8$. Theorem 6.4 yields:

Corollary 6.8. *The $\mathcal{G}^{E_6}$-module $\widehat{V(\lambda_3)}$ is irreducible if $c\in\mathbb{C}\setminus\{\mathbb{N}-15,-17,-19,-21\}$.*

Let k be a positive integer. We calculate by (6.35) that

$$\Upsilon(k\lambda_4)=\{(k+1)\lambda_4,(k+1)\lambda_4-\varepsilon_4+\varepsilon_5,(k-1)\lambda_4+\varepsilon_1\},\tag{6.112}$$

which yields $\ell_\omega(k\lambda_4)=k/2-6$. The fifth fundamental weight of $o(10,\mathbb{C})$ is $\lambda_5=(1/2)\sum_{i=1}^5\varepsilon_i$. Now (6.35) implies

$$\Upsilon(k\lambda_5)=\{k\lambda_5+\lambda_4,k\lambda_5+\lambda_4-\varepsilon_3-\varepsilon_4,(k-1)\lambda_5\},\tag{6.113}$$

which implies $\ell_\omega(k\lambda_5)=-k/2-10$.

We define a representation of $o(10,\mathbb{C})$ on $\mathcal{C}=\mathbb{C}[z_1,...,z_{16}]$ obtained from (2.49)-(2.71) with $\mathcal{A}$ replaced by $\mathcal{C}$ and x_i replaced by z_i for $i\in\overline{1,16}$. Then the $o(10,\mathbb{C})$-submodule $\mathcal{N}_k$ generated by z_1^k is isomorphic to $V(k\lambda_4)$. Note that by (6.67) and (6.68),

$$\Upsilon'(k\lambda_4)=\{k\lambda_4+\varepsilon_1,k\lambda_4+\varepsilon_5\}.\tag{6.114}$$

Note that $\zeta_1z_1^k$ is an $o(10,\mathbb{C})$-singular vector of weight $k\lambda_4+\varepsilon_1$ in $U\mathcal{N}_k$, where $M=\mathcal{N}_k$ in the earlier settings. By (2.71) with x_i replaced by z_i, (6.55) and (6.66),

$$T_1'(z_1^k)=\frac{k}{2}\zeta_1z_1^k\Longrightarrow\flat_{k\lambda_4+\varepsilon_1}=\frac{k}{2}.\tag{6.115}$$

By (2.49)-(2.53), (2.69) and (2.70) with x_i replaced by z_i,

$$(E_{9,10}-E_{5,4})(z_1^k)=kz_1^{k-1}z_2\in\mathcal{N}_k\Longrightarrow z_1^{k-1}z_2\in\mathcal{N}_k,\tag{6.116}$$

$$(E_{8,9}-E_{4,3})(z_1^{k-1}z_2)=z_1^{k-1}z_3\in\mathcal{N}_k,\tag{6.117}$$

$$(E_{3,2}-E_{7,8})(z_1^{k-1}z_3)=z_1^{k-1}z_4\in\mathcal{N}_k,\tag{6.118}$$

$$(E_{2,1}-E_{6,7})(z_1^{k-1}z_4)=z_1^{k-1}z_7\in\mathcal{N}_k.\tag{6.119}$$

According to (2.49)-(2.53) with x_i replaced by z_i and Table 1, we find that the vector

$$u'=\zeta_5z_1^k+\zeta_4z_1^{k-1}z_2+\zeta_3z_1^{k-1}z_3-\zeta_2z_1^{k-1}z_4+\zeta_1z_1^{k-1}z_7\tag{6.120}$$

is an $o(10,\mathbb{C})$-singular vector of weight $k\lambda_4+\varepsilon_5$ in $U\mathcal{N}_k$. Expressions (2.49)-(2.71) with x_i replaced by z_i, (6.55), (6.57)-(6.59) and (6.66),

$$\begin{aligned}&T_1'(z_1^{k-1}z_7)-T_2'(z_1^{k-1}z_4)+T_3'(z_1^{k-1}z_3)+T_4'(z_1^{k-1}z_2)+T_5'(z_1^k)\\ =\;&(k/2-1)\zeta_1z_1^{k-1}z_7+\zeta_2z_1^{k-1}z_4-\zeta_3z_1^{k-1}z_3-\zeta_4z_1^{k-1}z_2-\zeta_5z_1^k\\ &-\zeta_1z_1^{k-1}z_7-(k/2-1)\zeta_2z_1^{k-1}z_4-\zeta_3z_1^{k-1}z_3-\zeta_4z_1^{k-1}z_2-\zeta_5z_1^k\\ &-\zeta_1z_1^{k-1}z_7+\zeta_2z_1^{k-1}z_4+(k/2-1)\zeta_3z_1^{k-1}z_3-\zeta_4z_1^{k-1}z_2-\zeta_5z_1^k\\ &-\zeta_1z_1^{k-1}z_7+\zeta_2z_1^{k-1}z_4-\zeta_3z_1^{k-1}z_3+(k/2-1)\zeta_4z_1^{k-1}z_2-\zeta_5z_1^k\\ &-k\zeta_1z_1^{k-1}z_7+k\zeta_2z_1^{k-1}z_4-k\zeta_3z_1^{k-1}z_3-k\zeta_4z_1^{k-1}z_2-(k/2)\zeta_5z_1^k\\ =\;&-(k/2+4)u',\end{aligned}\tag{6.121}$$

equivalently, $\flat_{k\lambda_4+\varepsilon_5} = -(k/2+4)$. Thus $\flat(k\lambda_4) = -(k/2+4)$. Symmetrically, $\flat(k\lambda_5) = -(k/2+4)$. By Theorem 6.4, we have:

Corollary 6.9. *The* $\mathcal{G}^{E_6}$*-module* $\widehat{V(k\lambda_4)}$ *is irreducible if* $c \in \mathbb{C} \setminus \{\mathbb{N} - 10 - k/2, 2\mathbb{N} + k - 12\}$. *The* $\mathcal{G}^{E_6}$*-module* $\widehat{V(k\lambda_5)}$ *is irreducible if* $c \in \mathbb{C} \setminus \{\mathbb{N} - 10 - k/2, 2\mathbb{N} - k - 20\}$.

References

[**A**] J. Adams, *Lectures on Exceptional Lie Groups*, The University of Chicago Press Ltd., London, 1996.

[**AB1**] G. Anderson and T. Blažek, E_6 unification model building.I. Clebsch-Gordan coefficients of $27 \otimes \overline{27}$, *J. Math. Phys.* **41** (2000), no. 7, 4808-4816.

[**AB2**] G. Anderson and T. Blažek, E_6 unification model building.II. Clebsch-Gordan coefficients of $78 \otimes 78$, *J. Math. Phys.* **41** (2000), no. 12, 8170-8189.

[**AB3**] G. Anderson and T. Blažek, E_6 unification model building.III. Clebsch-Gordan coefficients in E_6 tensor products of the 27 with higher-dimensional representations, *J. Math. Phys.* **46** (2005), no. 1, 013506, 13pp.

[**As1**] M. Aschbacher, The 27-dimensional module for E_6.I., *Invent. Math.* **89** (1987), no. 1, 159-195.

[**BCDH**] P. Berglund, P. Candelas, X. de le Ossa, E. Derrick, J. Distler and T. Hübsch, On instanton contrbutions to the masses and couplings of E_6 singles, *Nuclear Phys. B* **454** (1995), no. 1-2, 127-163.

[**BZ**] B. Bineger and R. Zierau, A singular representation of E_6, *Trans. Amer. Math. Soc.* **341** (1994), no. 2, 771-785.

[**B-N**] J. Bion-Nadal, Subfactor of the hyperfinite Π_1 factor with Coxeter graph E_6 as invariant, *J. Operator Theory* **28** (1992), 27-50.

[**BK**] R. Brylinski and B. Kostant, Minimal representations of E_6, E_7, and E_8 and the generalized Capelli identity, *Proc. Nar. Acad. Sci. U.S.A.* **91** (1994), no. 7., 2469-2472.

[**CD**] B. Cerchiai and A. Scotti, Mapping the geometry of the E_6 group, *J. Math. Phys.* **49** (2008), no. 1, 012107, 19pp.

[DL] C. Das and L. Laperashvili, Preon model and family replicated E_6 unification, *SIGMA* **4** (2008), 012, 15pages.

[D] L. Dickson, A class of groups in an arbitrary realm connected with the configuration of the 27 lines on a cubic surface, *J. Math.* **33** (1901), 145-123.

[FGP] J. Fernández-Núãez, W. Garcia-Fuertes and A. Perelomov, Irreducible characters and Clebsch-Gordan series for the exceptional algebra E_6: an approach through the quantum Calogero-Sutherland model, *J. Nonlinear Math. Phys.* **12**, suppl. 1, 280-301.

[Gm] M. D. Gould, Tensor operators and projection techniques in infinite dimensional representations of semi-simple Lie algebras, *J. Phys. A: Math. Gen.* **17** (1984), 1-17.

[GSA] A. Ghezelbash, A. Shafiekhani and M. Abolbasani, On the Picard-Fuchs equations of $N = 2$ supersymmetric E_6 Yang-Mills theory, *Modern Phys. Lett. A* **13** (1998), no. 7, 527-531.

[G] D. Ginzburg, On standard L-functions for E_6 and E_7, *J. Reine Angew. Math.* **465** (1995), 101-131.

[HM] N. Haba and T. Matsuoka, Large lepton flavor mixing and E_6-type unification models, *Progr. Theoret. Phys.* **99** (1998), no. 5, 831-842.

[HH] J. E. M. Homos and Y. M. M. Homos, Algebraic model for the evolution of the generic code, *Phys. Rev. Lett.* **71** (1991), 4401-4404.

[Hub] R. Hubert, The (A_2, G_2) duality in E_6, octonions and the triality principle, *Trans. Amer. Math. Soc.* **360** (2008), no. 1, 347-367.

[Hum] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag New York Inc., 1972.

[HK] R. Howl and S. King, Minimal E_6 supersymmetric standard model, *J. High Ener. Phys.* **01**(2008), 030, 31pp.

[I] A. Iltyakov, On rational invariants of the group E_6, *Proc. Math. Soc.* **124** (1996), no. 12, 3637-3640.

[Ka] V. Kac, *Infinite-Dimensional Lie Algebras*, Birkhäser, Boston, Inc., 1982.

[K] B. Kostant, On the tensor product of a finite and an infinite dimensional representation, *J. Func. Anal.* **20** (1975), 257-285.

[MPW] I. Morrison, P. Pieruschka and B. Wybourne, The interacting boson model with the exceptional groups G_2 and E_6, *J. Math. Phys.* **32** (1991), no. 2, 356-372.

[OM] Y. Okamoto and R. Marshak, A garnd unification preon model with E_6 metacolor, *Nuclear Phys. B* **268** (1986), no. 2, 397-405.

[R] H. Rubenthaler, The (A_2, G_2) duality in E_6, octonians and the triality principle, *Trans. Amer. Math. Soc.* **360** (2008), no. 1, 347-367.

[S1] G. Shen, Graded modules of graded Lie algebras of Cartan type (I)—mixed product of modules, *Science in China A* **29** (1986), 570-581.

[S2] G. Shen, Graded modules of graded Lie algebras of Cartan type (II)—positive and negative graded modules, *Science in China A* **29** (1986), 1009-1019.

[S3] G. Shen, Graded modules of graded Lie algebras of Cartan type (III)—irreducible modules, *Chin. Ann. of Math B* **9** (1988), 404-417.

[SW] K. Suzuki and M. Wakui, On the Turaev-Viro-Ocneanu invariant of 3-manifolds derived from the E_6-subfactor, *Kyushu J. Math.* **56** (2002), 59-81.

[T] J. Tits, A local approach to buildings, in: C. Davis, B. Gronbaum and F. Sherk (eds), "Geometric Vein," Berlin-Heidelberg-New York, Springer, 1982, pp. 519-547.

[Wa] X. Wang, Identification of Gepner's model with twisted LG model and E_6 singlets, *Modern Phy. Lett. A* **6** (1991), no. 23, 2155-2162.

[X1] X. Xu, *Kac-Moody Algebras and Their Representations*, China Science Press, 2007.

[X2] X. Xu, A cubic E_6-generalization of the classical theorem on harmonic polynomials, *J. Lie Theory* **21** (2011), 145-164.

[X3] X. Xu, Representations of Lie algebras and coding theory, *J. Lie Theory*, accepted.

[XZ] X. Xu and Y. Zhao, Generalized conformal representations of orthogonal Lie algebras, *arXiv:1105.1254v1[math.RT].*

[ZX] Y. Zhao and X. Xu, Generalized projective representations for sl(n+1), *J. Algebra* **328** (2011), 132-154.